

EXISTENCE AND NON-EXISTENCE OF CONSTANT SCALAR CURVATURE AND EXTREMAL SASAKI METRICS

CHARLES P. BOYER, HONGNIAN HUANG, EVELINE LEGENDRE,
AND CHRISTINA W. TØNNESEN-FRIEDMAN

ABSTRACT. We discuss the existence and non-existence of constant scalar curvature, as well as extremal, Sasaki metrics. We prove that the natural Sasaki-Boothby-Wang manifold over the admissible projective bundles over local products of non-negative CSC Kähler metrics, as described in [ACGTF08a], always has a constant scalar curvature (CSC) Sasaki metric in its Sasaki-Reeb cone. Moreover, we give examples that show that the extremal Sasaki-Reeb cone, defined as the set of Sasaki-Reeb vector fields admitting a compatible extremal Sasaki metric, is not necessarily connected in the Sasaki-Reeb cone, and it can be empty even in the non-Gorenstein case. We also show by example that a non-empty extremal Sasaki-Reeb cone need not contain a (CSC) Sasaki metric which answers a question posed in [BHLTF21]. The paper also contains an appendix where we explore the existence of Kähler metrics of constant weighted scalar curvature, as defined in [Lah19], on admissible manifolds over local products of non-negative CSC Kähler metrics.

CONTENTS

1. Introduction	2
2. Extremal Sasaki geometry	3
2.1. Brief review on extremal Sasaki structures	3
2.2. The weighted extremal case	5
2.3. Admissible Kähler manifolds	6
3. Existence and non-existence	7
3.1. Sufficient conditions for CSC Sasaki metrics in the Sasaki-Reeb cone	7
3.2. The Sasaki-Futaki invariant in the admissible case	14
3.3. Sasaki-Reeb cones with extremal but no CSC Sasaki metrics	16
3.4. Sasaki-Reeb cones with no extremal Sasaki metrics	20
4. The topology of (M, \mathcal{S})	21
Appendix A. Constant weighted scalar curvature on admissible Kähler manifolds	23
A.1. Limitation of Theorem A.1	27
References	28

Date: October 25, 2022.

The first author was partially supported by grant #519432 from the Simons Foundation. The third author is partially supported by France ANR project BRIDGES No ANR-21-CE40-0017. The fourth author was partially supported by grant #422410 from the Simons Foundation.

1. INTRODUCTION

In the recent years, powerful tools have been developed to investigate the question of existence of extremal Kähler and Sasaki metrics, particularly regarding the more famous cases of Einstein and constant scalar curvature Kähler/Sasaki metrics. As expected, these tools, including various notions of stability [CS18] and weighted Kähler metrics [AC21, ACL21], are now used to explore the extremal problem within families of Kähler/Sasaki structures. This is the idea pursued in the present paper where we use weighted Kähler metrics together with admissible constructions on projective bundles to describe more precisely some *extremal Sasaki–Reeb cones*.

This cone is the Sasakian counterpart of the extremal Kähler cone (i.e, on a given a complex manifold, the cone of Kähler classes containing extremal Kähler metrics) but sits in the Lie algebra \mathfrak{t} of a compact torus T acting on complex cone (Y, \widehat{J}) . More precisely, for a given T -invariant complex cone (Y, \widehat{J}) , the Sasaki–Reeb cone $\mathfrak{t}^+(Y, \widehat{J})$ of (Y, \widehat{J}) , is the set of polarizations $\xi \in \mathfrak{t}$ admitting a compatible T -invariant radial Kähler potential $r \in C^\infty(Y, \mathbb{R}^+)$, that is r is proper, T -invariant and satisfies $\mathcal{L}_{-\widehat{J}\xi}r = r$, $dd^c r > 0$. The Sasaki–Reeb cone admits an equivalent definition in terms of CR-structure induces on the level set of each radial potential as we recall in §2.1.

The extremal Sasaki–Reeb cone, say $\mathfrak{e}(Y, \widehat{J})$, is then the subset of polarizations $\xi \in \mathfrak{t}^+(Y, \widehat{J})$ admitting a compatible T -invariant radial Kähler potential $r \in C^\infty(Y, \mathbb{R}^+)$ whose associated Sasaki metric on the level set $\{r = 1\}$ is extremal in the sense of Boyer–Galicki–Simanca [BGS08]. For example, if (Y, \widehat{J}) is the cone over a polarized Kähler manifold (X, L) admitting an extremal Kähler metric in the class $2\pi c_1(L)$ then the regular Sasaki–Reeb vector field (given by the fiberwise \mathbb{C}^* action) lies in $\mathfrak{e}(Y, \widehat{J})$.

Although the Sasaki–Reeb cone is known to be open, strictly convex and polyhedral [Ler02, BGS08, CS18, BvC18, ACL21], very little is known about the extremal Sasaki–Reeb cone. As proved in [BGS08], transversal scaling of Sasaki structures implies that $\mathfrak{e}(Y, \widehat{J})$ is a subcone of $\mathfrak{t}^+(Y, \widehat{J})$. Moreover, $\mathfrak{e}(Y, \widehat{J})$ is open, and can be empty or all of $\mathfrak{t}^+(Y, \widehat{J})$.

One contribution of the present article, is the exhibition of two families of examples of complex cones for which the extremal Sasaki–Reeb cone has interesting properties.

- **Examples 3.7** *non-Gorenstein examples where the extremal Sasaki–Reeb cone is empty; such examples were previously known only in the Gorenstein case [BvC18];*
- **Examples 3.5** *showing that extremal Sasaki–Reeb cone need not be connected.*

We have a special interest in constant scalar curvature Sasaki structures (CSCS). We denote $\mathfrak{csc}(Y, \widehat{J})$ the set of $\xi \in \mathfrak{t}^+(Y, \widehat{J})$ for which there exists a compatible T -invariant radial Kähler potential $r \in C^\infty(Y, \mathbb{R}^+)$ whose associated Sasaki metric on the level set $\{r = 1\}$ has constant scalar curvature. It is known, [BGS08, FOW09] that

$$(1) \quad \mathfrak{csc}(Y, \widehat{J}) = \mathfrak{e}(Y, \widehat{J}) \cap \{\xi \in \mathfrak{t}^+(Y, \widehat{J}) \mid \text{Fut}_\xi = 0\}$$

where Fut_ξ is the transversal (or Sasaki) Futaki invariant. Note that $\{\xi \in \mathfrak{t}^+(Y, \widehat{J}) \mid \text{Fut}_\xi = 0\}$ is non-empty for any complex cone over a compact Sasaki manifold, see [BHL18]. However, we provide an example here for which the intersection (1) is empty.

- **Examples** of Sasaki-Reeb cones with compatible extremal Sasaki metrics (i.e., $\mathfrak{e}(Y, \widehat{J}) \neq \emptyset$) but no compatible CSCS metrics (i.e., $\mathfrak{csc}(Y, \widehat{J}) = \emptyset$). See Theorem 3.12 and Example 3.5.

This last family of examples answers a question posed in [BHLTF21].

All the examples above are built in the context of the Sasakian geometry of Boothby-Wang manifolds (M, \mathcal{S}) over admissible Kähler manifolds as defined in [ACGTF08a], denoted here by N^{ad} . In particular, our main result is the following condition ensuring that the CSCS cone is non-empty for some of these important cases.

Theorem 1.1. *[see Theorem 3.1] Let Ω be an admissible Hodge Kähler class on the admissible projective bundle $N^{\text{ad}} = \mathbb{P}(E_0 \oplus E_\infty) \rightarrow N$, where N is a compact Kähler manifold which is a local product of nonnegative CSMK metrics. Then for the complex cone (Y, \widehat{J}) over (N^{ad}, Ω) , the Sasaki-Reeb cone $\mathfrak{t}^+(Y, \widehat{J})$ (relative to the 2-dimensional torus obtained from the double tower of fibrations) has a non-empty subset $\mathfrak{csc}(Y, \widehat{J})$ of compatible radial Kähler potentials with associated CSCS metrics.*

In previous work [BTF14, BTF16, BTF22c, BTF22b] the first and last authors have given many examples proving the existence of constant scalar curvature Sasaki metrics. The method of proofs used the admissible construction of projective bundles developed in [ACGTF08a] together with the Sasaki join construction [BGO07]. This latter construction, which can be thought of as the Sasaki analogue of de Rham decomposition of Kähler manifolds, divides Sasaki manifolds into two types, those which can be described as joins are said to be *decomposable*; whereas, those that cannot be described by joins are *indecomposable*. For a detailed description see [BHLTF18].

In the rest of the paper, the examples and results are stated in terms of CR structures and Sasaki metrics. In particular, Theorem 3.1 below is the Sasaki version of Theorem 1.1. Subsection 2.1, and in particular Remark 2.1, makes the transition between the CR-Sasaki and the complex cone dictionaries.

The key novel idea in the proof of Theorem 1.1 (Theorem 3.1) is to employ the link, discovered by [AC21], between the weighted extremal Kähler metrics and Sasaki extremal metrics and then use the properness of the Einstein-Hilbert functional [BHL18] to discover CSCS solutions in the admissible setting.

Acknowledgements. The authors thank Vestislav Apostolov and Ivan Cheltsov for helpful and encouraging correspondence.

2. EXTREMAL SASAKI GEOMETRY

2.1. Brief review on extremal Sasaki structures. To explain the notation and point of view we are using in this paper, let recall that a Sasakian structure on a manifold M^{2n+1} is a special type of contact metric structure $\mathcal{S} = (\xi, \eta, \Phi, g)$ with underlying almost CR structure (\mathcal{D}, J) where η is a contact form such that $\mathcal{D} = \ker \eta$,

ξ is its Reeb vector field, $J = \Phi|_{\mathcal{D}}$, and $g = d\eta \circ (\mathbb{1} \times \Phi) + \eta \otimes \eta$ is a Riemannian metric. \mathcal{S} is a Sasakian structure if ξ is a Killing vector field, so we call it a Sasaki-Reeb vector field, and the almost CR structure is integrable, i.e. (\mathcal{D}, J) is a CR structure. We refer to [BG08] for the fundamentals of Sasaki geometry. We call (\mathcal{D}, J) a *CR structure of Sasaki type*, and \mathcal{D} a *contact structure of Sasaki type*. We shall always assume that the Sasaki manifold M^{2n+1} is compact and connected.

Let $\mathcal{S} = (\xi, \eta, \Phi, g)$ be a T -invariant Sasakian structure where T is a compact torus acting effectively on M , that is $T \subset \text{Diff}(M)$ and such that $\xi \in \text{Lie } T =: \mathfrak{t}$. In this setting, the Sasaki–Reeb cone of (M, \mathcal{S}) , relatively to T , is

$$\mathfrak{t}^+(\mathcal{D}, J) := \{\tilde{\xi} \in \mathfrak{t} \mid \eta(\tilde{\xi}) > 0\}.$$

Each $\tilde{\xi} \in \mathfrak{t}^+(\mathcal{D}, J)$ determines a T -invariant Sasakian structure, compatible with (\mathcal{D}, J) with contact form $\eta(\tilde{\xi})^{-1}\eta$. Therefore, sometimes in this paper, and in the literature, elements of $\mathfrak{t}^+(\mathcal{D}, J)$ are called Sasakian metrics.

If instead of fixing a contact structure, we fix a Sasaki–Reeb vector field ξ and vary the contact structure, we obtain a contractable space of Sasakian structures (all compatible with ξ), namely

$$(2) \quad \mathcal{S}(\xi, \bar{J}) = \{\eta + d_B^c \varphi \mid \varphi \in C_B^\infty(M), (\eta + d_B^c \varphi) \wedge (d\eta + dd_B^c \varphi)^n \neq 0\},$$

where $d_B^c \varphi = -d\varphi \circ \Phi$. The space $\mathcal{S}(\xi, \bar{J})$ is an infinite dimensional Fréchet manifold. When starting with a T -invariant Sasakian structure as above, it is convenient to consider only the space of T -invariant isotopic contact structures $\mathcal{S}(\xi, \bar{J})^T$. Note that the Sasaki–Reeb cones, relative to T , of elements in $\mathcal{S}(\xi, \bar{J})^T$ all agree.

All the contact structures in $\mathcal{S}(\xi, \bar{J})$ have the same basic cohomology class $[d\eta]_B \in H^{1,1}(\mathcal{F}_\xi)$ and are compatible with the same holomorphic transversal structure (ξ, \bar{J}) . Each representative $\eta \in \mathcal{S}(\xi, \bar{J})$ determines a transverse Kähler structure with transverse Kähler metric $g^T = d\eta \circ (\mathbb{1} \times \Phi)$. Note that $d\eta$ is not exact as a basic cohomology class, since η is not a basic 1-form. We want to search for a ‘preferred’ Sasakian structure \mathcal{S}_φ which represents the basic cohomology class $[d\eta]_B$. This leads to the study [BGS08] of the Calabi functional

$$(3) \quad \mathcal{E}_2(g) = \int_M s_g^2 dv_g$$

where the variation is taken over the space $\mathcal{S}(\xi, \bar{J})$. As in the Kähler case the Euler–Lagrange equation is a 4th order PDE whose critical points are those Sasaki metrics whose transversal scalar curvature s_g is a *Killing potential*, that is, its gradient is transversely holomorphic, or, equivalently, the contact vector field of s_g preserves the CR structure. Such Sasaki metrics (structures) are called *extremal*. Note that a T -invariant Sasakian structure $\mathcal{S} = (\xi, \eta, \Phi, g)$ with a contact moment map $\mu : M \rightarrow \mathfrak{t}^*$, is extremal if s_g is the pull-back by μ of an affine-linear function on \mathfrak{t}^* . In practice, this is a convenient way to check that a Sasakian structure is extremal.

In this article we study the following set, that we call the *extremal Sasaki–Reeb cone*

$$\mathfrak{e}(\mathcal{D}, J) := \{\xi \in \mathfrak{t}^+(\mathcal{D}, J) \mid \exists \text{ an extremal Sasakian structure in } \mathcal{S}(\xi, \bar{J})^T\}.$$

Remark 2.1. To recover the definition of the extremal Sasaki–Reeb cone given in the introduction, recall from [ACL21] that $\mathcal{S}(\xi, \bar{J})^T$ is in one to one correspondence with the space of T -invariant radial potentials, relative to ξ , on the complex cone

(Y, \widehat{J}) over M . The complex structure \widehat{J} on Y depends on ξ and on the transversal holomorphic structure (ξ, \bar{J}) . However, given two commuting Sasaki–Reeb vector fields $\xi, \tilde{\xi} \in \mathfrak{t}^+(\mathcal{D}, J)$, their associated complex cones are biholomorphic [HS16]. Therefore, $\mathfrak{t}^+(\mathcal{D}, J) = \mathfrak{t}^+(Y, \widehat{J})$ and $\mathfrak{e}(\mathcal{D}, J) = \mathfrak{e}(Y, \widehat{J})$ where we identify a Reeb vector field ξ with its lift to (Y, \widehat{J}) .

An important special case of extremal Sasaki structures are those of constant scalar curvature (cscS) in which case the gradient of s_g is the zero vector field. An extremal Sasakian structure is CSC if and only if the transversal Futaki invariant $Fut_{\mathcal{S}} = Fut_{\xi} : \mathfrak{h}^T(\xi, \bar{J}) \rightarrow \mathbb{R}$ vanishes. Here $\mathfrak{h}^T(\xi, \bar{J})$ is the Lie algebra of transversally holomorphic vector fields, and we use the alternative definition (see eg. [BHLTF17])

$$Fut_{\xi}(X) = \int_M (s_g - \bar{s}_g) \eta(X) \eta \wedge (d\eta)^n$$

with $\bar{s}_g := \int_M s_g \eta(X) \eta \wedge (d\eta)^n / \int_M \eta \wedge (d\eta)^n$. Note that Fut_{ξ} depends only on the isotopy class $\mathfrak{S}(\xi, \bar{J})$ and not on the element used to represent it.

Thus, as described in the introduction, we get nested cones

$$\mathbf{csc}(\mathcal{D}, J) \subset \mathfrak{e}(\mathcal{D}, J) \subset \mathfrak{t}^+(\mathcal{D}, J)$$

which are the main topics of this paper.

2.2. The weighted extremal case. One effective way of treating the indecomposable case alluded to in the introduction is to use the weighted extremal construction developed in [AC21, ACL21, AJL]. It can also be considered as a generalization of [BTF22a] where the projective bundle is a twist one stage 3 Bott orbifold. Here we briefly review the weighted extremal approach. Let (N, g, ω) be a Kähler manifold of complex dimension m , f a positive Killing potential on N , and (weight) $p \in \mathbb{R}$. Then the (f, p) -Scalar curvature of g is given by

$$(4) \quad Scal_{f,p}(g) = f^2 Scal(g) - 2(p-1)f\Delta_g f - p(p-1)|df|_g^2,$$

If $Scal_{f,p}(g)$ is a Killing potential, g is said to be a (f, p) -extremal Kähler metric. The case $p = 2m$ has been studied by several people and is interesting due to the fact that $Scal_{f,2m}(g)$ computes the scalar curvature of the Hermitian metric $h = f^{-2}g$. The case of interest to us here is when $p = m + 2$. This case is related to the study of extremal Sasaki metrics [AC21, ACL21, AJL]. Indeed, if we assume that the Kähler class $[\omega/2\pi]$ is an integer class, then $\frac{Scal_{f,m+2}(g)}{f}$ is equal to the transverse scalar curvature of a certain Sasaki structure (determined by f) in the Sasaki-Reeb cone of the Boothby-Wang constructed Sasaki manifold over (N, g, ω) . More precisely, if χ is the Reeb vector field of the Sasaki structure coming directly from the Boothby-Wang construction over (N, g, ω) and f is viewed as a pull-back to the Sasaki manifold, then $\xi := f\chi$ is a Reeb vector field in the Sasaki-Reeb cone giving a new Sasaki structure. While the pull-back from N of $Scal(g)$ is the Tanaka-Webster scalar curvature of the Tanaka-Webster connection induced by χ , the expression $\frac{Scal_{f,m+2}(g)}{f}$ pulls back from N to be the Tanaka-Webster scalar curvature of the Tanaka-Webster connection induced by ξ . The latter is then also identified with the transverse scalar curvature of the Sasaki structure defined by ξ . This fact is seen from the details of the proof of Lemma 3 in [AC21]. As also

follows from Theorem 1 of [AC21], the Sasaki structure determined by f is extremal if and only if g is $(f, m+2)$ -extremal. In this case, the corresponding Sasaki structure is CSC if and only if $\frac{Scal_{f, m+2}(g)}{f}$ is constant.

2.3. Admissible Kähler manifolds. In this paper we study the Sasakian geometry of Boothby-Wang manifolds over the admissible Kähler manifolds defined in [ACGTF08a]. Admissible Kähler manifolds are very tractable for exploring the existence of Kähler metrics with special geometries. The original case in point is Calabi's construction of extremal Kähler metrics on Hirzebruch surfaces and related higher dimensional ruled manifolds [Cal82]. Subsequently these types of constructions have been explored by many (e.g. Guan [Gua95], Hwang [Hwa94], Hwang-Singer [HS02], Koiso [Koi90], Koiso-Sakane [KS86], LeBrun [LeB91], Pedersen-Poon [PP98], Simanca [Sim91], and Tønnesen-Friedman [TF98]). For full details for these manifolds we refer to e.g. [ACGTF08a] or Sections 2.1 and 2.2 of [AMTF]. Here we will just give a brief introduction:

Definition 2.2. *Let N be a compact Kähler manifold that is covered by a product of simply connected Kähler manifolds $(N_a, \pm\omega_a, \pm g_a)$ of dimensions d_a , $a \in \mathcal{A} = \{1, \dots, I\}$, and let E_0 and E_∞ be projectively flat Hermitian holomorphic vector bundles on N of ranks $d_0 + 1$ and $d_\infty + 1$, respectively, which satisfy the condition*

$$(*) \quad \frac{c_1(E_\infty)}{d_\infty + 1} - \frac{c_1(E_0)}{d_0 + 1} = \frac{1}{2\pi} \sum_{a \in \mathcal{A}} [\omega_a].$$

*Then the projective bundle $N^{ad} = \mathbb{P}(E_0 \oplus E_\infty) \rightarrow N$ is said to be **admissible**.*

Note that the complex dimension of N^{ad} is $m = \sum_{a \in \hat{\mathcal{A}}} d_a + 1$. Note also that the Equation (*) puts topological constraints on the manifolds N^{ad} . In particular, an admissible bundle is never trivial.

The fibers of such projective bundles admit an S^1 action which gives rise to (a canonical Kähler metric and) a moment map construction $\mathfrak{z} : N_0^{ad} \rightarrow (-1, 1)$ as described in [HS02, ACGTF08a] where $N_0^{ad} \subset N^{ad}$ is the subset of stable points of N^{ad} . This gives rise to a family of Kähler metrics which are defined on all of N^{ad} and locally described on N_0^{ad} by

$$(5) \quad g = \sum_{a \in \hat{\mathcal{A}}} \frac{1 + x_a \mathfrak{z}}{x_a} g_a + \frac{d\mathfrak{z}^2}{\Theta(\mathfrak{z})} + \Theta(\mathfrak{z})\theta^2, \quad \omega = \sum_{a \in \hat{\mathcal{A}}} \frac{1 + x_a \mathfrak{z}}{x_a} \omega_a + d\mathfrak{z} \wedge \theta,$$

where θ is a connection 1-form, Θ is a smooth function with domain containing $(-1, 1)$, and x_a for $a \in \mathcal{A}$, are real numbers of the same sign as g_a and satisfying $0 < |x_a| < 1$. Here $\hat{\mathcal{A}}$ is the finite set $\mathcal{A} = \{1, \dots, N\}$ if $d_0 = d_\infty = 0$, and $\hat{\mathcal{A}}$ equals $\mathcal{A} \cup \{0\}$ if $d_0 > 0, d_\infty = 0$, equals $\mathcal{A} \cup \{\infty\}$ if $d_0 = 0, d_\infty > 0$, and equals $\mathcal{A} \cup \{0\} \cup \{\infty\}$ if $d_0 > 0, d_\infty > 0$. Further, for $d_0 \neq 0$, we set $x_0 = 1$ and let g_0 be the Fubini-Study metric on $\mathbb{C}\mathbb{P}^{d_0}$ with scalar curvature $2d_0(d_0 + 1)$ (or $2d_0 s_0$ with $s_0 = d_0 + 1$). Similarly, for $d_\infty \neq 0$, we set $x_\infty = -1$ and let $-g_\infty$ be the Fubini-Study metric on $\mathbb{C}\mathbb{P}^{d_\infty}$ with scalar curvature $2d_\infty(d_\infty + 1)$ (or $-2d_\infty s_\infty$ with $s_\infty = -(d_\infty + 1)$). Moreover, Θ satisfies the following conditions:

$$(6) \quad (i) \Theta(\mathfrak{z}) > 0, \quad -1 < \mathfrak{z} < 1, \quad (ii) \Theta(\pm 1) = 0, \quad (iii) \Theta'(\pm 1) = \mp 2.$$

The triple (N^{ad}, ω, g) where N^{ad} is given in Definition 2.2 and (ω, g) in Equation (5), is called an **admissible Kähler manifold**.

With all else being fixed, the **admissible Kähler class** $\Omega = [\omega]$ is determined by the choice of x_a for $a \in \mathcal{A}$. If the x_a values are all rational, then, up to scale, Ω is a rational cohomology class. In this case, and for convenience, we simply say that Ω is rational. Note that a rational admissible Kähler class can be rescaled to be an integer class. In general, any rescale of an admissible Kähler class will also be called admissible.

3. EXISTENCE AND NON-EXISTENCE

In addition to the brief introduction above, we refer the reader to [ACGTF08a] (or the more recent summary in Sections 2.1 and 2.2 of [AMTF]) for a detailed description of admissible manifolds, their admissible metrics, and admissible Kähler classes. We also refer to these sources for the notation used below. Except for a couple of exceptions (pointed out below), we are staying faithful to the notation used in [ACGTF08a] and [AMTF]. In particular, we have a moment map $\mathfrak{z} : N^{ad} \rightarrow [-1, 1]$ (denoted by z in [ACGTF08a] and [AMTF]) associated to a natural S^1 -action of the admissible Kähler forms. Now, for the case where N is a local product of *non-negative* CSCK metrics, by Proposition 11 of [ACGTF08a] and (the second half of) Theorem 1 in [AMTF] we have that every admissible Kähler class admits an admissible $(c\mathfrak{z} + 1, p)$ -extremal metric for any choice of $c \in (-1, 1)$ and $p \in \mathbb{R}$. In more detail, when $c = 0$ an $(c\mathfrak{z} + 1, p)$ -extremal metric is just an extremal metric and we appeal to Proposition 11 of [ACGTF08a]. When $0 < |c| < 1$, by rescaling we can let $f = |\mathfrak{z} + \mathbf{a}|$, where $\mathbf{a} = 1/c$ and appeal to (the second half of) Theorem 1 in [AMTF]¹. Alternatively one can recast Section 2.3 as well as the proof of Theorem 3.1 in [AMTF], replacing $f = |z + \mathbf{a}|$ with $f = c\mathfrak{z} + 1$ and reprove the above existence result in a self-contained manner. For simplicity, we shall not summarize this whole process, but only extract exactly what we need for our purposes here.

3.1. Sufficient conditions for CSC Sasaki metrics in the Sasaki-Reeb cone.

Here we will assume that the admissible Kähler class Ω is rational and so there is a Boothby-Wang constructed Sasaki manifold (M, \mathcal{S}) given by an appropriate re-scale of Ω . The (re-scales of) Reeb vector fields induced by the Killing potentials $c\mathfrak{z} + 1$, for $c \in (-1, 1)$, span a 2-dimensional subcone of the (unreduced) Sasaki-Reeb cone. We note that the Sasaki-Reeb cone $\mathfrak{t}^+(\mathcal{S})$ of (M, \mathcal{S}) has dimension $d_0 + d_\infty + 2 + \dim(\mathfrak{t})$ where \mathfrak{t} is the Lie algebra of the maximal torus of the automorphism group $\mathfrak{Aut}(N)$ of the Kähler manifold N . We also note that $c = 0$ corresponds to the initial ray (since a constant Killing potential f corresponds to a rescale of the initial Reeb vector field coming from the Boothby-Wang construction). We are ready for

Theorem 3.1. *Suppose Ω is a rational admissible Kähler class on the admissible manifold $N^{ad} = \mathbb{P}(E_0 \oplus E_\infty) \rightarrow N$, where N is a compact Kähler manifold which is a local product of nonnegative CSCK metrics. Let (M, \mathcal{S}) be the Boothby-Wang constructed Sasaki manifold given by an appropriate rescale of Ω . Then the corresponding Sasaki-Reeb cone will always have a (possibly irregular) CSC-ray (up to isotopy).*

¹Note that the assumption of $\mathbf{a} > 1$ (and so $|\mathfrak{z} + \mathbf{a}| = \mathfrak{z} + \mathbf{a}$) in [AMTF] is merely practical and all the arguments are easily adapted to include the $\mathbf{a} < -1$ (and $|\mathfrak{z} + \mathbf{a}| = -\mathfrak{z} - \mathbf{a}$) case as well.

Proof. If the (admissible) extremal metric in Ω is CSC, we are done. In general we will show that there exist some $c \in (-1, 1)$ such that the extremal Sasaki structure determined by $f = c\mathfrak{z} + 1$ has constant scalar curvature.

Note that the admissible $(c\mathfrak{z} + 1, p)$ -extremal metric has $(c\mathfrak{z} + 1, p)$ -Scalar curvature $Scal_{c\mathfrak{z}+1,p} = A_1\mathfrak{z} + A_2$, where, adapting the details of Section 2.3 of [AMTF], A_1 and A_2 are given by the unique solution to the linear system

$$(7) \quad \begin{aligned} \alpha_{1,-(1+p)}A_1 + \alpha_{0,-(1+p)}A_2 &= 2\beta_{0,(1-p)} \\ \alpha_{2,-(1+p)}A_1 + \alpha_{1,-(1+p)}A_2 &= 2\beta_{1,(1-p)}, \end{aligned}$$

where

$$(8) \quad \begin{aligned} \alpha_{r,k} &= \int_{-1}^1 (ct + 1)^k t^r p_c(t) dt, \quad p_c(t) = \prod_{a \in \hat{A}} (1 + x_a t)^{d_a}, \\ \beta_{r,k} &= \int_{-1}^1 \left(\sum_{a \in \hat{A}} \frac{x_a d_a s_a}{1 + x_a t} \right) p_c(t) t^r (ct + 1)^k dt \\ &\quad + \left((-1)^r (1 - c)^k p_c(-1) + (1 + c)^k p_c(1) \right), \end{aligned}$$

where $s_a = \pm \frac{Scal_{\pm g_a}}{2d_a}$ is the normalized scalar curvature. Notice that

- $c\alpha_{r+1,k} + \alpha_{r,k} = \alpha_{r,k+1}$
- $c\beta_{r+1,k} + \beta_{r,k} = \beta_{r,k+1}$.

Now, for $p = m + 2$, where $m = \sum_{a \in \hat{A}} d_a + 1$, the extremal Sasaki structure determined by $f = c\mathfrak{z} + 1$ has constant scalar curvature if and only if $Scal_{c\mathfrak{z}+1,p} = A_1\mathfrak{z} + A_2$ is a constant multiple of $c\mathfrak{z} + 1$, i.e., if and only if $A_1 - cA_2 = 0$. Now, $A_1 - cA_2 = 0$ may be re-written as

$$(\alpha_{1,-(1+p)}\beta_{0,(1-p)} - \alpha_{0,-(1+p)}\beta_{1,(1-p)}) - c(\alpha_{1,-(1+p)}\beta_{1,(1-p)} - \alpha_{2,-(1+p)}\beta_{0,(1-p)}) = 0,$$

which in turn is equivalent to $\alpha_{1,-p}\beta_{0,(1-p)} - \alpha_{0,-p}\beta_{1,(1-p)} = 0$, which with $p = m + 2$ is

$$(9) \quad \alpha_{1,-(m+2)}\beta_{0,-(m+1)} - \alpha_{0,-(m+2)}\beta_{1,-(m+1)} = 0.$$

Let K_c denote the Reeb vector field in the Sasaki-Reeb cone of \mathcal{S} induced by $f_{K_c} = c\mathfrak{z} + 1$. That is, $\eta_{\mathfrak{D}}^X(K_c) = c\mathfrak{z} + 1$ (when \mathfrak{z} is lifted to \mathcal{S} and X is the initial Reeb vector field of the Boothby-Wang construction alluded to in Theorem 3.1). In Section 3.2 we shall see directly that Equation (9) is an obstruction to the vanishing of the Sasaki-Futaki invariant Fut_{K_c} . Inspired by this fact and Lemma 3.1 in [BHLTF17], we turn our attention to the *Einstein-Hilbert functional*:

Up to an overall positive rescale, the Einstein-Hilbert functional introduced in Section 3 of [BHLTF17] is given by

$$H_S(K_c) := H_S(c) = S_c^{m+1}/V_c^m,$$

where $V_c = \alpha_{0,-(m+1)}$ and

$$S_c = \int_{-1}^1 Scal_{c\mathfrak{z}+1,m+2}(g)(c\mathfrak{z} + 1)^{-(m+2)} p_c(\mathfrak{z}) d\mathfrak{z} = 2\beta_{0,-m}.$$

Thus, up to an overall positive rescale, $H_S(c) = \frac{(\beta_{0,-m})^{m+1}}{(\alpha_{0,-(m+1)})^m}$. Using that

- $\frac{d}{dc} [\alpha_{0,-(m+1)}] = -(m+1)\alpha_{1,-(m+2)}$
- $\frac{d}{dc} [\beta_{0,-m}] = -m\beta_{1,-(m+1)}$,

we easily verify that

$$\begin{aligned} H'_S(c) &= \frac{m(m+1)(\beta_{0,-m})^m(\alpha_{0,-(m+1)})^{m-1}}{(\alpha_{0,-(m+1)})^{2m}} [\alpha_{1,-(m+2)}\beta_{0,-m} - \alpha_{0,-(m+1)}\beta_{1,-(m+1)}] \\ &= \frac{m(m+1)(\beta_{0,-m})^m(\alpha_{0,-(m+1)})^{m-1}}{(\alpha_{0,-(m+1)})^{2m}} [\alpha_{1,-(m+2)}\beta_{0,-(m+1)} - \alpha_{0,-(m+2)}\beta_{1,-(m+1)}]. \end{aligned}$$

Now $\alpha_{0,-(m+1)} > 0$ and $\beta_{0,-m} > 0$ (the latter is due to the fact that we are assuming N is a local product of *nonnegative* CSCK metrics) and therefore critical points of $H_S(c)$ correspond exactly to solutions of Equation (9) which in turn would give us that an extremal Sasaki structure determined by $f = c\mathfrak{z} + 1$ that has constant scalar curvature. Showing that $H_S(c)$ must have a least one critical point $c \in (-1, 1)$, would finish the proof.

To that end, first notice that $H_S(c) = \frac{(\beta_{0,-m})^{m+1}}{(\alpha_{0,-(m+1)})^m}$ is a smooth positive function for $c \in (-1, 1)$. To show that $H_S(c)$ has a critical point inside $(-1, 1)$, we will show that $\lim_{c \rightarrow \pm 1^\mp} H_S(c) = +\infty$. Since $c \rightarrow \pm 1^\mp$ represents K_c approaching the boundary of the Sasaki cone, this actually follows directly from Lemma 5.1 in [BHL18], but for the sake of transparency we will verify this directly.

First notice that

$$\alpha_{0,-(m+1)} = \int_{-1}^1 (ct+1)^{-(m+1)}(1+t)^{d_0} p_0(t) dt = \int_{-1}^1 (ct+1)^{-(m+1)}(1-t)^{d_\infty} p_\infty(t) dt,$$

where $p_0(t) = p_c(t)/(1+t)^{d_0}$ and $p_\infty(t) = p_c(t)/(1-t)^{d_\infty}$ are polynomials of degree $m-1-d_0$ and $m-1-d_\infty$, respectively. Keep in mind that the complex dimension $m = \sum_{a \in \hat{\mathcal{A}}} d_a + 1$ (with $d_a > 0$ for $a \in \mathcal{A}$ and $d_a \geq 0$ for $a \in \hat{\mathcal{A}}$) and note that $p_0(-1) > 0$ and $p_\infty(1) > 0$. From the definition of $p_c(t)$ in Equation (8) we also know that $p_0(t)$ and $p_\infty(t)$ are positive for $-1 < t < 1$. By repeated integration by parts, we can see that $\alpha_{0,-(m+1)}$ is a rational function of c which is positive and has no vertical asymptotes within the interval $(-1, 1)$. To understand the behavior near $c = \pm 1$, we consider repeated integration by parts and observe that

$$\alpha_{0,-(m+1)} = \frac{\left(\frac{(m-1-d_0)! d_0! p_0(-1)}{m! c^{d_0+1}} + o(1-c) \right)}{(1-c)^{m-d_0}} = \frac{\left(\frac{(-1)^{d_\infty+1} (m-1-d_\infty)! d_\infty! p_\infty(1)}{m! c^{d_\infty+1}} + o(1+c) \right)}{(1+c)^{m-d_\infty}},$$

where $o(x)$ is a function which is smooth in a neighborhood of $x = 0$ and satisfies that $o(0) = 0$. In summary, this means that

$$\alpha_{0,-(m+1)} = \frac{\delta_0(c) + o(1-c)}{(1-c)^{m-d_0}} = \frac{\delta_\infty(c) + o(1+c)}{(1+c)^{m-d_\infty}},$$

where

$$\lim_{c \rightarrow 1} \delta_0(c) > 0 \quad \lim_{c \rightarrow -1} \delta_\infty(c) > 0.$$

Similarly, we can show, albeit a bit messier, that $\beta_{0,-(m+1)}$ is a rational function of c which is non-negative, has no vertical asymptotes within the interval $(-1, 1)$ and satisfies

$$\beta_{0,-m} = \frac{\gamma_0(c) + o(1-c)}{(1-c)^{m-d_0}} = \frac{\gamma_\infty(c) + o(1+c)}{(1+c)^{m-d_\infty}},$$

where

$$\lim_{c \rightarrow 1} \gamma_0(c) > 0 \quad \lim_{c \rightarrow -1} \gamma_\infty(c) > 0.$$

We refer to proof of the generalized statement in Lemma A.3 for more detail. Putting this together we have

$$H_S(c) = \frac{(\gamma_0(c) + o(1-c))^{m+1}}{(\delta_0(c) + o(1-c))^m} (1-c)^{-(m-d_0)} = \frac{(\gamma_\infty(c) + o(1+c))^{m+1}}{(\delta_\infty(c) + o(1+c))^m} (1+c)^{-(m-d_\infty)},$$

and hence clearly $\lim_{c \rightarrow \pm 1^\mp} H_S(c) = +\infty$. This together with the fact that $H_S(c)$ is smooth and bounded from below over the interval $(-1, 1)$ allows us to conclude that $H_S(c)$ has at least one critical point $c \in (-1, 1)$. For this value of c , we have $H'_S(c) = 0$ and therefore c solves (9) making K_c a CSC Sasaki Reeb vector field. If c is not a rational number, the corresponding CSC Sasaki structure will be irregular. \square

Remark 3.2. It follows from the proof of Theorem 3.1 that the CSC Sasaki metric lies in the 2-dimensional subcone of $\mathfrak{t}^+(\mathcal{S})$ spanned by the constants plus $c\mathfrak{z} + 1$. Note that here we view the Sasaki-Reeb cone in terms of Killing potentials.

Example 3.1. As a simple example we let $N = \mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1$ and $N_{\mathbf{k}}$ be the total space of the twist 1 stage 3 Bott tower $\mathbb{P}(\mathbb{1} \oplus \mathcal{O}(k_1, k_2)) \rightarrow \mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1$ as studied in [BCTF19]. If $k_1 k_2 > 0$ the Kähler manifold $N_{\mathbf{k}}$ does not have a CSCCK metric. Nevertheless, Theorem 3.1 says that there is a CSCS ray in the Sasaki cone of the Boothby-Wang constructed Sasaki 7-manifold M^7 over $N_{\mathbf{k}}$. This also follows from the Main Theorem in [BTF22a]. The CSCCK obstruction here is the well known non-reductivity obstruction of Matsushima and Lichenerowicz. The topology of these 7-manifolds M^7 was described in [BTF22a] where they were shown to have the rational cohomology of the 2-fold connected sum $2\#(S^2 \times S^5)$ with torsion in H^4 which generally depends on both the pair (k_1, k_2) and the integral Kähler class on $N_{\mathbf{k}}$. We refer to Section 3 of [BTF22a] for details.

Example 3.2. Let N be a projective K3 surface. Then N has a non-trivial holomorphic line bundle L , so the $\mathbb{C}\mathbb{P}^1$ bundle $N^{ad} = \mathbb{P}(\mathbb{1} \oplus L)$ over N is also non-trivial. Moreover, Theorem 7 of [ACGTF08a] implies that the projective bundle N^{ad} does not admit a CSC Kähler metric. But again, Theorem 3.1 says that there is a CSCS ray in the Sasaki cone of the Boothby-Wang constructed Sasaki 7-manifold M^7 over N^{ad} . Thus, choosing an admissible (up to scale) primitive Kähler class on N^{ad} gives a simply connected M^7 with $H^2(M^7, \mathbb{Z}) \approx \mathbb{Z}^{22}$ with a CSC Sasakian structure which may be irregular. Such 7-manifolds can be constructed as nontrivial S^3 bundles over K3 surfaces by using the fiber join construction of Yamazaki [Yam99, BTF21]. Let us describe a bit more of the geometry and topology of such M^7 . This gives manifolds of the form

$$S^3 \longrightarrow M^7 \xrightarrow{\pi} N, \quad M^7 = M_1^5 \star_f M_2^5$$

where M_i^5 are Sasaki 5-manifolds formed from the Boothby-Wang construction over a projective K3 surface N with integral Kähler form ω_i . By Theorem 10.3.8 of [BG08] M_1^5 and M_2^5 are both diffeomorphic to the connected sum $21\#(S^2 \times S^3)$. Moreover, as described in [BTF21] the 7-manifolds M^7 divide into two types depending on whether the classes $[\omega_1], [\omega_2]$ are colinear or non-colinear. From Proposition 3.12 of [BTF21] and the fact that K3 has trivial canonical bundle, we see that the first Chern class of the contact bundle \mathcal{D} on M^7 satisfies $c_1(\mathcal{D}) = -\pi^*([\omega_1] + [\omega_2])$. In the colinear case we have $\omega_i = l_i\omega$ where ω is primitive Kähler form. In this case we see that the second Stiefel-Whitney class $w_2(M^7)$ vanishes if and only if $l_1 + l_2$ is even; hence, M^7 is spin when $l_1 + l_2$ is even and non-spin when $l_1 + l_2$ is odd. In both cases the 4-skeleton is formal, so for each such 3-sphere bundle it follows from Theorem 2.2 of [KT91] that there is a finite number of diffeomorphism types. Thus, since c_1 is a contact invariant, in both cases there is a diffeomorphism class on M^7 that has a countable infinity of inequivalent contact structures of Sasaki type.

Combining Theorem 3.1 with Corollary 1.1 of [CS18] gives the following statement concerning K-stability.

Corollary 3.3. *Under the hypothesis of Theorem 3.1 there exists a K-semistable ray of Sasakian structures in the Sasaki-Reeb cone.*

Remark 3.4. Note that Theorem 1 of [AC21] (the proof of which is local in nature) also holds for Sasaki orbifolds. Further, for $d_0 = d_\infty = 0$ (the so-called *no-blow-down case*), replacing each “ $p_c(-1)$ ” by “ $p_c(-1)/m_\infty$ ” and each “ $p_c(1)$ ” by “ $p_c(-1)/m_0$,” where $m_0, m_\infty \in \mathbb{Z}^+$, in the proof of Theorem 3.1, corresponds to generalizing the endpoint condition (iii) in (6) from

$$\Theta'(-1) = 2 \quad \text{and} \quad \Theta'(1) = -2.$$

to

$$\Theta'(-1) = 2/m_\infty \quad \text{and} \quad \Theta'(1) = -2/m_0.$$

Since this generalization does not affect the core of the proof, we can see that Theorem 3.1 generalizes within the no-blow-down case to the case of admissible metrics with orbifold singularities along the zero (D_0) and infinity (D_∞) sections of $N^{ad} = \mathbb{P}(E_0 \oplus E_\infty) \rightarrow N$. In other words N^{ad} can be replaced by the log pair (N^{ad}, Δ) , where $\Delta = (1 - 1/m_0)D_0 + (1 - 1/m_\infty)D_\infty$.

Suppose now that the admissible manifold in Theorem 3.1 is Fano, i.e. $c_1(N^{ad}) > 0$. Consider a Boothby-Wang constructed Sasaki manifold over N^{ad} given by some Kähler form representing $c_1(N^{ad})/\mathcal{J}$, where \mathcal{J} denotes the index of N^{ad} . Then $c_1(\mathcal{D}) = 0$ and so the CSC Sasaki metrics from Theorem 3.1 above are now η -Einstein. Second, we see from Lemma 5.2/Proposition 5.3 in [BHL18] that since the basic first Chern class of the initial Sasaki metric is positive (as a pullback of the positive class $c_1(N^{ad})/\mathcal{J}$), the average transverse scalar curvature of any Sasaki structure in the Sasaki-Reeb cone must be positive. In particular, the transverse (constant) scalar curvature of any η -Einstein structure in the cone must be positive. This means that any η -Einstein ray admits a Sasaki-Einstein structure. Thus, in this case the corresponding Sasaki-Reeb cone will always have a (possibly irregular) Sasaki-Einstein structure (up to isotopy). Therefore, in the Gorenstein case we have the following corollary which in turn is a special case of

the existence results by T. Mabuchi and Y. Nakagawa [MN13] as well as V. Apostolov, S. Jubert, and A. Lahdili (see e.g. Theorem 3 of [AJL]).

Corollary 3.5. *Suppose N^{ad} is a Fano admissible manifold $N^{ad} = P(E_0 \oplus E_\infty) \rightarrow N$, where N is a compact Kähler manifold which is a product of positive KE metrics. Let (M, \mathcal{S}) be the Boothby-Wang constructed Sasaki manifold given by $c_1(N^{ad})/\mathcal{J}$. Then the corresponding Sasaki-Reeb cone will always have a (possibly irregular) Sasaki-Einstein metric (up to isotopy).*

Example 3.3. For a simple non-toric example using the above corollary, let us assume that $\mathcal{A} = \{1\}$ and $N = N_1$ is the smooth Fermat sextic del Pezzo threefold $V_1 = z_0^6 + z_1^6 + z_2^6 + z_3^3 + z_4^2 = 0$ in $\mathbb{C}\mathbb{P}(\mathbf{w}) = \mathbb{C}\mathbb{P}(1, 1, 1, 2, 3)$ described in Example 3.4.2. of [ACC⁺21] and [CS09]. We have $|\mathbf{w}| - d = 2$ so this is Fano with index 2. It is K-stable and admits a KE metric. Furthermore, since there is only one quadratic term the automorphism group is finite (see e.g. Proposition 37 in [BGK05]) implying that it is a non-toric KE manifold. We also assume that g_1 is the KE metric on V_1 with $s_1 = 2$ and that $d_0 = 0$. Then, independent of the value of $d_\infty \in \mathbb{N} \cup \{0\}$, the projective bundle $N^{ad} = \mathbb{P}(\mathbb{1} \oplus E_\infty) \rightarrow N$ is Fano² and the Kähler class $c_1(N^{ad})/\mathcal{J}$ is a positive scalar multiple of the admissible class determined by $x_1 = \frac{d_\infty + 2}{d_\infty + 4}$ (see equations (3.8) and (3.11) in [ACGTF08b]). One can now calculate directly that for this value of x_1 and $c = 0$, the left hand side of equation (9) equals $\frac{-3 \cdot 2^{2d_\infty + 11} (2d_\infty + 5)(11d_\infty + 28)}{(d_\infty + 1)(d_\infty + 4)^8 (d_\infty + 5)} \neq 0$. This means that the Futaki invariant does not vanish for $c_1(N^{ad})/\mathcal{J}$, but Corollary 3.5 gives us a quasiregular or irregular SE metric in the Sasaki-Reeb cone of the $(2(4 + d_\infty) + 1)$ -dimensional, non-toric, Boothby-Wang constructed Sasaki manifold (M, \mathcal{S}_1) given by $c_1(N^{ad})/\mathcal{J}$.

Remark 3.6. In general, the Fano conditions for an admissible manifold $N^{ad} = \mathbb{P}(E_0 \oplus E_\infty) \rightarrow N$ are summarized in Theorem 3.1 of [ACGTF08b]. If we assume no-blow-down here we have $d_0 = d_\infty = 0$. Specifically, assume $N = \prod_{a \in \mathcal{A}} N_a$ is a finite product of compact positive Kähler-Einstein manifolds $(N_a, \pm g_a, \pm \omega_a)$ with scalar curvature $\pm 2d_a s_a$ and let $N^{ad} = \mathbb{P}(\mathbb{1} \oplus \mathcal{L}) \rightarrow N$, where $\mathcal{L} = \otimes_{a \in \mathcal{A}} \mathcal{L}_a$ and \mathcal{L}_a is (the pull-back of) a holomorphic line bundle over N_a such that $c_1(\mathcal{L}_a) = [\frac{\omega_a}{2\pi}]$. Then the Fano condition is exactly the condition that $s_a > 1$ if ω_a is positive and $s_a < -1$ if ω_a is negative. Since the Ricci forms $\rho_a = s_a \omega_a$, we see that a necessary condition for this to be a possible choice is that the index J_a of each N_a is at least 2. Further, from the details of Section 3 in [ACGTF08b] (specifically equation (3.8) in [ACGTF08b]) we observe that if $s_a \neq s_{\bar{a}}$, then $x_a \neq x_{\bar{a}}$, where the choice of x_a for each $a \in \mathcal{A}$ is governed by the Kähler class $c_1(N^{ad})/\mathcal{J}$ (which is then an appropriate rescale of an admissible Kähler class). Indeed, it is not hard to see that (3.8) in [ACGTF08b] for $d_0 = d_\infty = 0$ simply gives us $x_a = 1/s_a$.

Remark 3.7. Varying the complex structure on N gives rise to a bouquet of Sasaki-Reeb cones on M as described in [Boy13, BTF13, BTF14].

Example 3.4. Let us exhibit another example of a non-toric M satisfying the assumptions in Corollary 3.5 so that $c_1(N^{ad})/\mathcal{J}$ itself is not KE; hence, it is K-unstable. For

²If we instead assumed that $-g_1$ was KE and $s_1 = -2$, then the Fano condition would be that $d_\infty = 0$ whereas d_0 could be any value in $\mathbb{N} \cup \{0\}$.

simplicity we will assume we are in the no-blow-down case and that $N = N_1 \times N_2$ (so here $\hat{\mathcal{A}} = \mathcal{A} = \{1, 2\}$). Let N_1 be equal to $\mathbb{C}\mathbb{P}^2$ and pick $s_1 = 3/2$ (meaning ω_1 is 4π times the unit volume Fubini-Study Kähler form). Then $x_1 = 2/3$. Let N_2 be equal to the smooth Fermat sextic del Pezzo threefold $V_1 = z_0^6 + z_1^6 + z_2^6 + z_3^3 + z_4^2 = 0$ in $\mathbb{C}\mathbb{P}(\mathbf{w}) = \mathbb{C}\mathbb{P}(1, 1, 1, 2, 3)$ described in Example 3.3. We have $|\mathbf{w}| - d = 2$ so this is Fano with index 2. It is K-stable and admits a KE metric. Furthermore, since there is only one quadratic term the automorphism group is finite (see e.g. Proposition 37 in [BGK05]) implying that $N_1 \times N_2$ is non-toric KE manifold.

We now consider the projective bundle N^{ad} over $N_1 \times N_2$. Since the Fano index is 2, we can pick $s_2 = 2$. Then $x_2 = 1/2$. For the admissible data $(s_1, s_2, x_1, x_2, d_1, d_2) = (3/2, 2, 2/3, 1/2, 2, 3)$ and $c = 0$ we can easily check that (9) does not hold. Equivalently, plugging this admissible data into (7) of [ACGTF08a], we calculate that $\alpha_1\beta_0 - \alpha_0\beta_1 \neq 0$. According to Proposition 6 in [ACGTF08a] we therefore have that the Futaki invariant does not vanish for $c_1(N^{ad})/J$. This means that $c_1(N^{ad})/J$ itself is not CSC, hence not KE. Corollary 3.5 now gives us a quasiregular or irregular SE metric in the Sasaki-Reeb cone of the 13-dimensional, non-toric, Boothby-Wang constructed Sasaki manifold (M, \mathcal{S}_1) given by $c_1(N^{ad})/J$. Note that since $x_1 \neq x_2$, this example does not overlap with any of the examples arising from $S_{\mathbf{w}}^3$ -joins (see [BTF16, BTF22c]). Furthermore, varying the complex structure on V_1 induces a bouquet of Sasaki-Reeb cones on 13-manifold (M, \mathcal{S}_1) , and it follows from [AZ21] that each smooth complex structure admits a Sasaki-Einstein metric. Since V_1 has no continuous automorphisms, the dimension of the local moduli space is

$$h^0(\mathbb{C}\mathbb{P}(\mathbf{w}), \mathcal{O}(d)) - \sum_i h^0(\mathbb{C}\mathbb{P}(\mathbf{w}), \mathcal{O}(w_i)) =$$

$$h^0(\mathbb{C}\mathbb{P}(\mathbf{w}), \mathcal{O}(6)) - h^0(\mathbb{C}\mathbb{P}(\mathbf{w}), \mathcal{O}(1)) - h^0(\mathbb{C}\mathbb{P}(\mathbf{w}), \mathcal{O}(2)) - h^0(\mathbb{C}\mathbb{P}(\mathbf{w}), \mathcal{O}(3)).$$

We get $h^0(\mathbb{C}\mathbb{P}(\mathbf{w}), \mathcal{O}(6)) = 59$, $h^0(\mathbb{C}\mathbb{P}(\mathbf{w}), \mathcal{O}(1)) = 3$, $h^0(\mathbb{C}\mathbb{P}(\mathbf{w}), \mathcal{O}(2)) = 7$, and $h^0(\mathbb{C}\mathbb{P}(\mathbf{w}), \mathcal{O}(3)) = 13$ giving a 36-dimensional complex family. This gives rise to a bouquet of one dimensional Sasaki-Reeb cones on the S^1 bundle over V_1 corresponding to the primitive Kähler class which in turn induces a bouquet of Sasaki-Reeb cones on the 13-dimensional manifold (M, \mathcal{S}_1) .

It should be clear that many such examples can be constructed using results in [BGK05] as well as [ACC⁺21]. So we now consider a fairly large class \mathcal{C} of simply connected algebraic varieties generalizing our example. We consider Brieskorn-Pham polynomials of degree d with exponent vector $\mathbf{a} \in (\mathbb{Z}^+)^{n+1}$, weight vector $\mathbf{w} \in (\mathbb{Z}^+)^{n+1}$ of the form

$$(10) \quad f_{\mathbf{a}}(\mathbf{z}) = z_0^{a_0} + \cdots + z_n^{a_n}, \quad a_i = \frac{d}{w_i}.$$

We do not want linear terms in $f_{\mathbf{a}}$ since they give rise to the standard sphere. So we shall assume that all $a_j \geq 2$. There is a natural weighted \mathbb{C}^* action on \mathbb{C}^{n+1} given by

$$(11) \quad \lambda \cdot \mathbf{z} = (\lambda^{w_0} z_0, \dots, \lambda^{w_n} z_n)$$

which induces

$$f_{\mathbf{a}}(\lambda \cdot \mathbf{z}) = \lambda^d f_{\mathbf{a}}(\mathbf{z}).$$

So the Brieskorn manifold $M_{\mathbf{a}}$ defined by $M_{\mathbf{a}} = (f_{\mathbf{a}}(\mathbf{z}) = 0) \cap S^{2n+1}$ has a natural weighted S^1 action defined by restriction. The weighted \mathbb{C}^* quotient of the hypersurface $(f_{\mathbf{a}}(\mathbf{z}) = 0) \setminus \{0\}$ or equivalently the weighed S^1 quotient of $M_{\mathbf{a}}$ gives weighted projective varieties of the form

$$(12) \quad V_{\mathbf{a}} = \text{Proj } \mathbb{C}[z_0, \dots, z_n]/(f_{\mathbf{a}})$$

where (f) denotes the ideal generated by f . These varieties all embed in the weighted projective space $\mathbb{C}\mathbb{P}(\mathbf{w})$. Brieskorn manifolds $M_{\mathbf{a}}$ also have a natural Sasakian structure whose Reeb vector field ξ generate the S^1 action. In order to satisfy the hypothesis of Theorem 3.1 we want the S^1 quotient of $M_{\mathbf{a}}$ to be a smooth manifold with a trivial orbifold structure³. Equivalently we want the Sasakian structure on $M_{\mathbf{a}}$ to be regular (cf. Proposition 9.3.22 of [BG08]).

Proposition 3.8. *A Brieskorn manifold has a regular Sasakian structure $\mathcal{S}_{\mathbf{w}}$ if and only if the weights are pairwise relatively prime.*

We are interested in regular Sasaki-Brieskorn manifolds whose projective algebraic quotient is Fano. The latter condition means that the Sasakian structure is positive which for a Brieskorn manifold is the condition $|\mathbf{w}| - d > 0$ where $|\mathbf{w}| = \sum_j w_j$. Here we restrict ourselves to dimensions 5 and 7 which correspond to $n = 3, 4$. As previously, we assume that $a_j \geq 2$, so standard spheres are eliminated.

Proposition 3.9. *Let $M_{\mathbf{a}}$ be a regular Brieskorn manifold of dimension 5 or 7 with a positive Sasakian structure. Then for*

- (1) $n = 3$; $\mathbf{a} = (2, 2, 2, 2), (3, 3, 3, 3), (4, 4, 4, 2), (6, 6, 3, 2)$;
- (2) $n = 4$; $\mathbf{a} =$
 $(2, 2, 2, 2, 2), (3, 3, 3, 3, 3), (4, 4, 4, 4, 4), (4, 4, 4, 4, 2), (6, 6, 6, 6, 2), (6, 6, 6, 3, 2)$.

Proof. The conditions that we must satisfy are $|\mathbf{w}| - d > 0$, the weights w_j are pairwise relatively prime, and divide d with no w_j equal to d . We do the case $n = 4$. First $\mathbf{w} = (1, 1, 1, 1, 1)$ implies $d = 2, 3, 4$ giving the first three on the list. If $\mathbf{w} = (1, 1, 1, 1, b)$ then we must have $b + 4 > d = ab$ for some $a \geq 2$. This implies that $a = 2$ and $b = 2, 3$ giving the next two. Next if $\mathbf{w} = (1, 1, 1, b, c)$ with $b < c$ relatively prime, then we have $b + c + 3 > ac$ for some $a \geq 2$. So $b + 3 > c$ which implies that $c = b + 1$ or $b + 2$. This implies $\mathbf{w} = (1, 1, 1, b, b + 1)$ or $\mathbf{w} = (1, 1, 1, b, b + 2)$. The only solution that satisfies $|\mathbf{w}| > d$ is $\mathbf{w} = (1, 1, 1, 2, 3)$ with $d = 6$. One can check that there are no further solutions. \square

In particular, as long as the Fano index of the projective variety $V_{\mathbf{a}}$ of Equation (12), with \mathbf{a} given in Proposition 3.9, is at least $d_0 + 2$, we can take N in Corollary 3.5 to be $V_{\mathbf{a}}$.

3.2. The Sasaki-Futaki invariant in the admissible case. Assume that we are in the admissible case as described in Section 2 of [AMTF] and assume that the admissible Kähler class Ω is rational and is represented by an admissible metric g corresponding to a smooth function $F(\mathfrak{z})$ satisfying (16) of [AMTF]). After an appropriate (and otherwise

³There are such smooth projective varieties with non-trivial orbifold structures, but we do not consider them here.

ignored) rescaling, this defines a Boothby-Wang constructed Sasaki manifold (M, \mathcal{S}) with Reeb vector field X .

Let K_c denote the Reeb vector field in the Sasaki-Reeb cone of (M, \mathcal{S}) induced by $f_{K_c} = c\mathfrak{z} + 1$. That is, $\eta_{\mathfrak{D}}^X(K_c) = c\mathfrak{z} + b$ (when \mathfrak{z} is lifted to (M, \mathcal{S})). It now follows from Section 2 of [AMTF] that the weighted scalar curvature $Scal_{c\mathfrak{z}+1, m+2}(g)$ is given by

$$Scal_{c\mathfrak{z}+1, m+2}(g) = \frac{-(c\mathfrak{z} + 1)^{m+3}G''(\mathfrak{z})}{p_c(\mathfrak{z})} + 2(c\mathfrak{z} + 1)^2 \left(\sum_{a \in \hat{A}} \frac{x_a d_a s_a}{1 + x_a \mathfrak{z}} \right),$$

where $G(\mathfrak{z}) := \frac{F(\mathfrak{z})}{(c\mathfrak{z}+1)^{m+1}}$ and $m = (\sum_{a \in \hat{A}} d_a) + 1$.

Now, consider the Futaki Invariant $Fut_{K_c}(Z)$ of Definition B.1. of [ACL21] for any $Z \in \mathfrak{t}$. Using Remark B.2. of [ACL21], we have that, up to a multiple by a non-zero constant,

$$Fut_{K_c}(Z) = \int_{-1}^1 (Scal_{c\mathfrak{z}+1, m+2}(g) - C_{K_c}(c\mathfrak{z} + 1)) f_Z(c\mathfrak{z} + 1)^{-m-3} p_c(\mathfrak{z}) d\mathfrak{z},$$

where $f_Z = \eta_{\mathfrak{D}}^X(Z)$ and, using (48) of [ACL21],

$$C_{K_c} = \frac{\int_{-1}^1 Scal_{c\mathfrak{z}+1, m+2}(g)(c\mathfrak{z} + 1)^{-m-2} p_c(\mathfrak{z}) d\mathfrak{z}}{\int_{-1}^1 (c\mathfrak{z} + 1)^{-m-1} p_c(\mathfrak{z}) d\mathfrak{z}} = \frac{2\beta_{0, -m}}{\alpha_{0, -m-1}}.$$

Let $Z = \mathfrak{z}X$ with \mathfrak{z} lifted to (M, \mathcal{S}) . We can view Z as $\frac{d}{dc}K_c$. Using the useful observation from the proof of Theorem 3.1 that

- $c\alpha_{r+1, k} + \alpha_{r, k} = \alpha_{r, k+1}$
- $c\beta_{r+1, k} + \beta_{r, k} = \beta_{r, k+1}$,

it is straightforward to verify that

$$\begin{aligned} (13) \quad Fut_{K_c}(Z) &= \frac{2(\alpha_{0, -(m+1)}\beta_{1, -(m+1)} - \alpha_{1, -(m+2)}\beta_{0, -m})}{\alpha_{0, -(m+1)}} \\ &= \frac{2(\alpha_{1, -(m+2)}\beta_{0, -(m+1)} - \alpha_{0, -(m+2)}\beta_{1, -(m+1)})}{\alpha_{0, -(m+1)}} \\ &= \frac{2}{\alpha_{0, -(m+1)}} \text{—times the left hand side of (9).} \end{aligned}$$

Thus equation (9) is equivalent with the vanishing of $Fut_{K_c}(Z)$.

In conclusion, (9) is apparently an obstruction to the existence (up to isotopy) of a CSC Sasaki structure in the ray determined by K_c . Of course, this also follows from the more “expensive” Theorem 3 in [ACL21] and the discussion leading up to (9). Indeed, combining Theorem 3 in [ACL21] and Proposition 2.2 of [AMTF] with the above observations we can state a very explicit existence criterion. To do so we need a few preliminary notations lifted straight from [AMTF] with some adaptation for our choice of $f = c\mathfrak{z} + 1$:

Let A_1 and A_2 be given by the unique solution to the linear system (7) and let

$$Q(\mathfrak{z}) = \frac{p_c(\mathfrak{z})}{(c\mathfrak{z} + 1)^{m+1}} \left(\sum_{a \in \hat{A}} \frac{x_a d_a s_a}{1 + x_a \mathfrak{z}} \right) - \frac{(A_1 \mathfrak{z} + A_2) p_c(\mathfrak{z})}{(c\mathfrak{z} + 1)^{m+3}}.$$

Now let

$$G(\mathfrak{z}) = \frac{2p_c(-1)}{(1-c)^{m+1}}(\mathfrak{z} + 1) + \int_{-1}^{\mathfrak{z}} Q(t)(\mathfrak{z} - t) dt$$

and consider the *weighted extremal polynomial*

$$(14) \quad F_{\Omega, c, m+2}(\mathfrak{z}) = (c\mathfrak{z} + 1)^{m+1}G(\mathfrak{z}).$$

Proposition 3.10. *Let Ω be a rational admissible Kähler class on an admissible Kähler manifold $N^{ad} = \mathbb{P}(E_0 \oplus E_\infty) \rightarrow N$ where N is a compact Kähler manifold which is a local product of CSCK metrics. Pick any admissible Kähler metric g with Kähler form in Ω and let the moment map be given by $\mathfrak{z} : N^{ad} \rightarrow [-1, 1]$. Let (M, \mathcal{S}) be the Boothby-Wang constructed Sasaki manifold given by an appropriate rescale of Ω . Let $c \in (-1, 1)$ and consider the corresponding weighted extremal polynomial $F_{\Omega, c, m+2}(\mathfrak{z})$ given by (14). Then*

- [ACL21] *the Reeb vector field K_c determined by $f_{K_c} = c\mathfrak{z} + 1$ is extremal (up to isotopy) if and only if*

$$F_{\Omega, c, m+2}(\mathfrak{z}) > 0, \quad -1 < \mathfrak{z} < 1$$

- *the Reeb vector field K_c determined by $f_{K_c} = c\mathfrak{z} + 1$ is CSC (up to isotopy) if and only if*

$$F_{\Omega, c, m+2}(\mathfrak{z}) > 0, \quad -1 < \mathfrak{z} < 1, \quad \text{AND} \quad \underbrace{\alpha_{1, -m-2}\beta_{0, (-m-1)} - \alpha_{0, -m-2}\beta_{1, (-m-1)}}_{\text{Equation(9)}} = 0,$$

where $\alpha_{r, k}$ and $\beta_{r, k}$ are given by (8).

Remark 3.11. Note that $F_{\Omega, 0, m+2}(\mathfrak{z})$ is the usual admissible extremal polynomial for Ω (see Definition 1 in [ACGTF08a]).

3.3. Sasaki-Reeb cones with extremal but no CSC Sasaki metrics. Existence results like Theorem 3.1 give rise to the question of whether a Sasaki CR structure with extremal metrics always has a CSC Sasaki metric in its Sasaki-Reeb cone. This was formulated as Problem 6.1.2 in [BHLTF21]. Here we give a negative answer to this question, namely:

Theorem 3.12. *There exist Sasaki CR structures with extremal Sasaki metrics, but no CSC Sasaki metrics.*

Proof. In principle we are using that a Sasakian structure (M, \mathcal{S}) in the extremal subcone $\mathfrak{e}(\mathcal{D}, J) \subset \mathfrak{t}^+(\mathcal{D}, J)$ is CSC if and only if its Futaki invariant $\mathcal{F}_{\mathcal{S}}$ vanishes and we note that $\mathcal{F}_{\mathcal{S}}$ only depends on the isotopy class [BGS08]. The proof proceeds by constructing an appropriate Sasaki CR structure over a certain smooth projective algebraic variety as described by the following example. \square

Remark 3.13. Before presenting the example we note that Theorem 3.12 is false when restricted to Sasaki CR structures with a 1-dimensional Sasaki-Reeb cone.

Example 3.5. We begin by revisiting the example presented carefully in Section 6 of [MTF11]. The admissible projective variety in question is $N^{ad} = \mathbb{P}(\mathcal{O} \oplus \mathcal{O}(1, -1)) \rightarrow \Sigma_1 \times \Sigma_2$, where Σ_i are compact genus two Riemann surfaces and g_1 and $-g_2$ are Kähler metrics on Σ_1 and Σ_2 , respectively that have constant scalar curvature equal to -4 .

Following the notation in [ACGTF08a], this means that $s_1 = -2$, $s_2 = 2$, $0 < x_1 < 1$, and $-1 < x_2 < 0$, where the choice of $\mathbf{x} = (x_1, x_2)$ determines the Kähler class

$$\Omega_{\mathbf{x}} = \frac{[\omega_1]}{x_1} + \frac{[\omega_2]}{x_2} + \Xi,$$

where Ξ is the Poincare dual of $2\pi(e_0 + e_\infty)$. Since here $[\omega_i]$ and Ξ are primitive integer classes, we see that $\Omega_{\mathbf{x}}$ is 2π times a rational class if and only if x_i are rational numbers. Note that this example does not satisfy the conditions assumed in Theorem 3.1.

Now, as explained in Section 6 of [MTF11], when $x_1 = x$ and $x_2 = -x$ for $0 < x < 1$, the Futaki invariant of the corresponding Kähler class vanishes (so any extremal metric in that class would be CSC) and the class contains a CSC Kähler metric if and only if the *extremal polynomial*, $F_x(\mathfrak{z}) = \frac{(1-\mathfrak{z}^2)(6-7x^2-4x^3+x^4-x^2(1-4x-x^2)\mathfrak{z}^2)}{2(3-x^2)}$, is positive for $-1 < \mathfrak{z} < 1$. The latter follows from Theorem 1 in [ACGTF08a]. As also explained in [MTF11], for $0 < x < 1$ sufficiently small, this positivity holds and we have CSC Kähler metrics in the corresponding class (as expected from Theorem 1 of [ACGTF08a]), but when e.g. $x = 4/5$, positivity fails and the corresponding class has no CSC Kähler metric and no extremal Kähler metric at all. The class can still be represented by some admissible Kähler metric. Moving forward we assume $x_1 = x$, $x_2 = -x$, and $x = 4/5$. Note that $F_{4/5}(\mathfrak{z}) = \frac{(1-\mathfrak{z}^2)(568\mathfrak{z}^2-37)}{1475}$.

Consider a Boothby-Wang constructed Sasaki manifold (M, \mathcal{S}) above N^{ad} with respect to an appropriate rescale of the Kähler class $\Omega_{4/5}$ determined by $x_1 = 4/5$, $x_2 = -4/5$ (pick e.g. a (rescale of an) admissible Kähler metric in $\Omega_{4/5}$). Then the (rescales of) Reeb vector fields K_c induced by the Killing potentials $c\mathfrak{z}+1$ for $c \in (-1, 1)$ span the entire Sasaki-Reeb cone of (M, \mathcal{S}) (since genus of $\Sigma_i = 2$, this cone has dimension 2)).

Using (7) for this particular case (where $p_c(t) = (1 - (4/5)^2 t^2)$) with $p = 5$, we calculate that $A_1 = -\frac{4c(535c^4+11566c^2-17933)}{115c^4-3686c^2+4543}$ and $A_2 = \frac{2(12305c^6-13795c^4+6227c^2-16401)}{115c^4-3686c^2+4543}$. By Proposition 3.10, we see that a ray in the Sasaki-Reeb cone corresponding to a value $c \in (-1, 1)$ admits an extremal Sasaki metric (up to isotopy) if and only if

$$F_{\Omega_{4/5}, c, 5}(\mathfrak{z}) = \frac{(1-\mathfrak{z}^2)(115c^4(553\mathfrak{z}^2-535)+110070c^3\mathfrak{z}(1-\mathfrak{z}^2)+c^2(98651-164999\mathfrak{z}^2)-149760c\mathfrak{z}(1-\mathfrak{z}^2)+154(568\mathfrak{z}^2-37))}{50(115c^4-3686c^2+4543)}$$

is positive for $-1 < \mathfrak{z} < 1$.

As expected from Remark 3.11, $F_{\Omega_{4/5}, 0, 5}(\mathfrak{z}) = F_{4/5}(\mathfrak{z})$. So for $-1 < c < 1$ sufficient close to zero, this positivity certainly fails. On the other hand, one can easily verify that

$$F_{\Omega_{4/5}, -1, 5}(\mathfrak{z}) = \frac{1}{300}(1-\mathfrak{z}^2)(-245\mathfrak{z}^3-86\mathfrak{z}^2+245\mathfrak{z}+194)$$

and

$$F_{\Omega_{4/5}, 1, 5}(\mathfrak{z}) = \frac{1}{300}(1-\mathfrak{z}^2)(245\mathfrak{z}^3-86\mathfrak{z}^2-245\mathfrak{z}+194)$$

both are positive for $-1 < \mathfrak{z} < 1$. Thus we may conclude that for $c \in (-1, 1)$ sufficiently close to either -1 or $+1$, $F_{\Omega_{4/5}, c, 5}(\mathfrak{z})$ is positive for $-1 < \mathfrak{z} < 1$.

We also can calculate that in this case equation (9) is equivalent to

$$(15) \quad \frac{4c(-2461c^4+512c^2+3893)}{5625(1-c^2)^7} = 0.$$

Note that this equation has no solution for $0 < |c| < 1$.

From the above work we see that, while \mathcal{S} itself (“initial ray”) is not extremal, for $c \in (-1, 1)$ sufficiently close to either -1 or $+1$ (“far enough away from the initial ray”), the corresponding ray does have an extremal Sasaki metric (up to isotopy). Moreover, since here equation (9) (i.e. equation (15)) has no solution for all $0 < |c| < 1$, there are no CSC rays in the Sasaki-Reeb cone at all. [Essentially, the only shot at a CSC ray would have been the initial ray of the Boothby-Wang construction, but since this ray allows no extremal metric at all, it is not a CSC ray either.] This example proves Theorem 3.12 and tells us that the answer is “no” to the question in Problem 6.1.2. of [BHLTF21].

We can dig a little deeper and numerically investigate the extremal Sasaki cone $\mathfrak{e}(\mathcal{D}, J)$ in $\mathfrak{t}^+(\mathcal{D}, J)$. In this case we identify $\mathfrak{t}^+(\mathcal{D}, J)$ with the interval $\mathcal{C} := \{c \mid -1 < c < -1\}$ swept out by transverse homotheties, and then correspondingly $\mathfrak{e}(\mathcal{D}, J)$ is identified with

$$\mathcal{C}_\epsilon := \{c \in \mathcal{C} \mid \forall \mathfrak{z} \in (-1, 1), F_{\Omega_{4/5}, c, 5}(\mathfrak{z}) > 0\} = \{c \in \mathcal{C} \mid \forall \mathfrak{z} \in (-1, 1), P(\mathfrak{z}, c) > 0\},$$

swept out by transverse homotheties where


$$\begin{aligned} P(\mathfrak{z}, c) &= 115c^4(553\mathfrak{z}^2 - 535) + 110070c^3\mathfrak{z}(1 - \mathfrak{z}^2) + c^2(98651 - 164999\mathfrak{z}^2) \\ &- 149760c\mathfrak{z}(1 - \mathfrak{z}^2) + 154(568\mathfrak{z}^2 - 37) \\ &= -5698 + 98651c^2 - 61525c^4 - 149760c\mathfrak{z} + 110070c^3\mathfrak{z} + 87472\mathfrak{z}^2 \\ &- 164999c^2\mathfrak{z}^2 + 63595c^4\mathfrak{z}^2 + 149760c\mathfrak{z}^3 - 110070c^3\mathfrak{z}^3. \end{aligned}$$

Equivalently, $\mathcal{C} \setminus \mathcal{C}_\epsilon = \{c \in \mathcal{C} \mid \exists \mathfrak{z} \in (-1, 1) \text{ s.t. } P(\mathfrak{z}, c) \leq 0\}$. To understand the latter set we need to see what part of the graph of $y = P(\mathfrak{z}, c)$, as defined over the square $\{(\mathfrak{z}, c) \mid -1 < \mathfrak{z} < 1, -1 < c < 1\}$, is below or at “sea level” (i.e. at or below $y = 0$).

Rather than attempting a rigorous treatment here, we shall rely on Mathematica to give us some clue: The plot of $y = P(\mathfrak{z}, c)$ looks as follows:

```
In[4]:= P := -5698 + 98651 c^2 - 61525 c^4 - 149760 c z + 110070 c^3 z + 87472 z^2 - 164999 c^2 z^2 +
63595 c^4 z^2 + 149760 c z^3 - 110070 c^3 z^3
```


```
In[6]:= Plot3D[{P, 0}, {z, -1, 1}, {c, -1, 1}, Axes -> True, AxesLabel -> {z, c},
PlotLabels -> "Expressions"]
```



From this picture it looks like $\mathcal{C} \setminus \mathcal{C}_\epsilon$ is one connected subset of \mathcal{C} , splitting \mathcal{C}_ϵ into two connected pieces. This is further underscored by the fact that the plot of $P(\mathfrak{z}, c) = 0$

in the square $\{(z, c) \mid -1 < z < 1, -1 < c < 1\}$ appears to be a simple closed curve:

```
In[8]:= ContourPlot[P == 0, {z, -1, 1}, {c, -1, 1}, Axes -> True, AxesLabel -> {z, c}]
```



We are now interested in finding the maximum and minimum c -values on this curve. This should give an estimate of $\mathcal{C} \setminus \mathcal{C}_c$, which seems to be an interval. To that end, by the theory of Lagrange Multipliers together with the picture at hand, we need to solve the system

$$\frac{\partial P}{\partial z} = 0$$

$$P = 0$$

over the interval $\{(z, c) \mid -1 < z < 1, -1 < c < 1\}$. We calculate that

$$\frac{\partial P}{\partial z} = 2(-74880c + 55035c^3 + 87472z - 164999c^2z + 63595c^4z + 224640cz^2 - 165105c^3z^2)$$

and find that numerically the above system has two solutions in $\{(z, c) \mid -1 < z < 1, -1 < c < 1\}$ with c values approximately equal to ± 0.410752 :

```
In[12]:= Factor[D[P, z]]
```

```
Out[12]= 2 (-74 880 c + 55 035 c^3 + 87 472 z - 164 999 c^2 z + 63 595 c^4 z + 224 640 c z^2 - 165 105 c^3 z^2)
```

```
In[13]:= NSolve[
```

```
{P == 0, (-74 880 c + 55 035 c^3 + 87 472 z - 164 999 c^2 z + 63 595 c^4 z + 224 640 c z^2 - 165 105 c^3 z^2) == 0},
{z, c}]
```

```
Out[13]= {{z -> -1.26012, c -> -3.00809}, {z -> 1.26012, c -> 3.00809},
{z -> 1.25913, c -> 1.12022}, {z -> -1.25913, c -> -1.12022}, {z -> 0.520035, c -> 1.30575},
{z -> -0.520035, c -> -1.30575}, {z -> 0.400666 - 0.682489 i, c -> -0.116165 - 0.420975 i},
{z -> 0.400666 + 0.682489 i, c -> -0.116165 + 0.420975 i},
{z -> -0.400666 - 0.682489 i, c -> 0.116165 - 0.420975 i},
{z -> -0.400666 + 0.682489 i, c -> 0.116165 + 0.420975 i},
{z -> -0.878086 - 0.642373 i, c -> 1.19307 - 0.028707 i},
{z -> -0.878086 + 0.642373 i, c -> 1.19307 + 0.028707 i},
{z -> 0.878086 - 0.642373 i, c -> -1.19307 - 0.028707 i},
{z -> 0.878086 + 0.642373 i, c -> -1.19307 + 0.028707 i},
{z -> -0.311144, c -> -0.410752}, {z -> 0.311144, c -> 0.410752}}
```

In conclusion, *the numerical evidence gathered from Mathematica tells us that $\mathcal{C} \setminus \mathcal{C}_c = [-\hat{c}, \hat{c}]$ and $\mathcal{C}_c = (-1, -\hat{c}) \cup (\hat{c}, 1)$, where $\hat{c} \approx 0.410752$.*

Another example can be given using the Yamazaki fiber join construction.

Example 3.6. Note that if instead we have $N^{ad} = \mathbb{P}(\mathcal{O} \oplus \mathcal{O}(8k_2^1, -8k_1^2)) \rightarrow \Sigma_1 \times \Sigma_2$, where genus of Σ_1 is $1 + 8k_2^1$, and genus of Σ_2 is $1 + 8k_1^2$, we still have $s_1 = -2$ and $s_2 = 2$ as in Example 3.5. If we further use the Kähler class corresponding to $x_1 = -x_2 = 4/5$ on N^{ad} , we have the same calculations and conclusion as above. In this case, we can view N^{ad} with Ω as the regular quotient of a Yamazaki fiber join as described in Section 5.3 of [BTF21] with $K = \begin{pmatrix} 9k_2^1 & k_1^2 \\ k_2^1 & 9k_1^2 \end{pmatrix}$.

3.4. Sasaki-Reeb cones with no extremal Sasaki metrics. There are many examples of Gorenstein Sasaki-Reeb cones with no extremal metrics whatsoever [BvC18]. Here we provide examples in the non-Gorenstein case.

Example 3.7. We will assume $N^{ad} = \mathbb{P}(\mathcal{O} \oplus \mathcal{O}(1, 1)) \rightarrow \Sigma_1 \times \Sigma_2$, where Σ_i are compact Riemann surfaces of genus 101 and 51 respectively. Let g_1 and g_2 be Kähler metrics on Σ_1 and Σ_2 , with constant scalar curvature equal to -400 and -200 , respectively. This means that $s_1 = -200$, $s_2 = -100$, $0 < x_i < 1$. Again, using Proposition 3.10, we calculate that for a Boothby-Wang constructed Sasaki manifold with respect to an appropriate rescale of the rational Kähler class Ω determined by $x_1 = 100/101$ and $x_2 = 9/10$, respectively, the Reeb vector field K_c induced by the Killing potential $c\mathfrak{z} + 1$ for $c \in (-1, 1)$ is extremal if and only if

$$F_{\Omega, c, 5}(\mathfrak{z}) = \frac{(1 - \mathfrak{z}^2)g_c(\mathfrak{z})}{2020(9714895c^4 - 41234400c^3 + 65557582c^2 - 46274160c + 12236095)}$$

is positive for $-1 < \mathfrak{z} < 1$, where

$$\begin{aligned} g_c(\mathfrak{z}) &= -2(230646399215c^4 - 1016343615871c^3 + 1668631275785c^2 - 1210756275690c + 327823027600) \\ &+ (-338414479445c^4 + 1516758787660c^3 - 2526484662221c^2 + 1856228939280c - 508090145290)\mathfrak{z} \\ &+ 2(249201848665c^4 - 1095101319871c^3 + 1793846257405c^2 - 1299139921290c + 351193969050)\mathfrak{z}^2 \\ &+ (375505948555c^4 - 1674191726860c^3 + 2776783510297c^2 - 2032903682160c + 554807556000)\mathfrak{z}^3. \end{aligned}$$

Since

$$\begin{aligned} &9714895c^4 - 41234400c^3 + 65557582c^2 - 46274160c + 12236095 \\ &= 9714895(1 - c)^4 + 2374820(1 - c)^3 + 143752(1 - c)^2 + 2616(1 - c) + 12 \\ &> 0 \end{aligned}$$

for all $c \in (-1, 1)$, positivity of $F_{\Omega, c, 5}(\mathfrak{z})$ for $-1 < \mathfrak{z} < 1$ is equivalent to positivity of $g_c(\mathfrak{z})$ for $-1 < \mathfrak{z} < 1$. However, e.g.

$$g_c(0) = -2(230646399215c^4 - 1016343615871c^3 + 1668631275785c^2 - 1210756275690c + 327823027600)$$

is negative for all $c \in (-1, 1)$, so positivity of $F_{\Omega, c, 5}(\mathfrak{z})$ for $-1 < \mathfrak{z} < 1$ will not hold for any value of $c \in (-1, 1)$. Similarly to Example 3.5, the Sasaki-Reeb cone here is parametrized by $c \in (-1, 1)$ and thus it follows that, in this case, the Sasaki-Reeb cone has no extremal rays at all.

Example 3.8. Note that if we instead we have $N^{ad} = \mathbb{P}(\mathcal{O} \oplus \mathcal{O}(200k_2^1, 18k_2^2)) \rightarrow \Sigma_1 \times \Sigma_2$, where genus of Σ_1 is $20000k_2^1 + 1$, and genus of Σ_2 is $900k_2^2 + 1$, we still have $s_1 = -200$, $s_2 = -100$ as in Example 3.7. If we further use the Kähler class corresponding to $x_1 = 100/101$ and $x_2 = 9/10$ on N^{ad} , we have the same calculations and conclusion as above. In this case, we can view N^{ad} with Ω as the regular quotient of a Yamazaki fiber join as described in Section 5.3 of [BTF21] with $K = \begin{pmatrix} 201k_2^1 & 19k_2^2 \\ k_2^1 & k_2^2 \end{pmatrix}$. Thus there is a countably infinite family of Yamazaki fiber joins that have no extremal Sasaki metrics at all.

Remark 3.14. These non-Gorenstein examples all have a 2-dimensional Sasaki-Reeb cone \mathfrak{t}^+ , as opposed to the examples in [BvC18]) where \mathfrak{t}^+ can be bigger.

4. THE TOPOLOGY OF (M, \mathcal{S})

Here we describe, more generally, the topology of the Boothby-Wang Sasaki manifold over the projective bundle N^{ad} some of which has been described in previous work [BTF22a] as well as briefly discussed in examples 3.1 and 3.2.

Proposition 4.1. *Let $S^1 \rightarrow M \rightarrow N^{ad} = \mathbb{P}(E_0 \oplus E_\infty)$ be the Boothby-Wang-Sasaki bundle. Then M is the total space of a lens space bundle over N .*

Proof. We have the commutative diagram of oriented fiber bundles

$$(16) \quad \begin{array}{ccccc} & & & & \mathbb{C}\mathbb{P}^{d_0+d_\infty+1} \\ & & & & \downarrow \\ S^1 & \longrightarrow & M & \xrightarrow{\pi_M} & N^{ad} \\ & & \searrow & & \downarrow \pi_N \\ & & & & N \end{array}$$

We claim that the fibers of the southeast arrow are lens spaces. By definition the southeast arrow is the projection $\pi_N \circ \pi_M$ which is clearly smooth. Let $p \in N$, the fiber $\pi_N^{-1}(p)$ is the projective space $\mathbb{C}\mathbb{P}^{d_0+d_\infty+1}$, and $\pi_M^{-1}(\pi_N^{-1}(p))$ is an S^1 bundle over $\pi_N^{-1}(p) = \mathbb{C}\mathbb{P}^{d_0+d_\infty+1}$ which is classified by $H^2(\mathbb{C}\mathbb{P}^{d_0+d_\infty+1}, \mathbb{Z}) \approx \mathbb{Z}$. Since M has a Sasakian structure, the bundle is determined by some $l \in \mathbb{Z}^+$. So the fiber of $\pi_N \circ \pi_M$ is a lens space $S^{2d_0+2d_\infty+3}/\mathbb{Z}_l$ where l is determined by the restriction of the Kähler form ω on N^{ad} to the fibers. The action of \mathbb{Z}_l on $S^{2d_0+2d_\infty+3}$ is given by

$$(z_1, \dots, z_{d_0+d_\infty+2}) \mapsto (\lambda z_1, \dots, \lambda z_{d_0+d_\infty+2})$$

where λ is an l th root of unity. □

We have

Theorem 4.2. *Let M be Boothby-Wang bundle over an admissible projective bundle $N^{ad} = \mathbb{P}(E_0 \oplus E_\infty) \rightarrow N$. Then*

- (1) $\pi_1(N^{ad}) = \pi_1(N)$;
- (2) $\pi_1(M)$ is an extension of $\pi_1(N)$ by \mathbb{Z}_l ;

(3) there is a spectral sequence converging to $H^*(M, \mathbb{Z})$ with E_2 term

$$E_2^{p,q} = H^p(N, \mathcal{H}^q(S^{2d_0+2d_\infty+3}/\mathbb{Z}_l, \mathbb{Z}))$$

where \mathcal{H}^q is the derived functor sheaf. Furthermore, if N is simply connected then $E_2^{p,q} = H^p(N, H^q)$ where

$$H^q = \begin{cases} \mathbb{Z} & \text{if } q = 0 \\ \mathbb{Z}_l & \text{if } q = 2, \dots, 2d_0 + 2d_\infty + 2 \\ \mathbb{Z} & \text{if } q = 2d_0 + 2d_\infty + 3 \\ 0 & \text{otherwise} \end{cases}$$

Proof. Items (1) and (2) follow from the exact homotopy sequences of the bundles in Proposition 4.1 and its proof. Item (3) follows from the Leray spectral sequence of the oriented lens space bundles of Proposition 4.1 together with the well known cohomology of lens spaces [BT82]. \square

If we consider the cohomology with \mathbb{Q} coefficients and the dimension of N is less than $2d_0 + 2d_\infty + 4$, the spectral sequence collapses and we get

Corollary 4.3. *If $\dim_{\mathbb{C}} N < d_0 + d_\infty + 2$, the Sasaki manifold M has the rational cohomology groups of the product $S^{2d_0+2d_\infty+3} \times N$.*

Example 4.1 (continuation of Example 3.3). We want to describe the topology of our Sasaki manifold M in Example 3.3. We have

Lemma 4.4. *The cohomology of the 6-manifold $N = V_1$ is*

$$H^q(N, \mathbb{Z}) = \begin{cases} \mathbb{Z} & \text{if } q = 0, 2, 4, 6 \\ \mathbb{Z}^{42} & \text{if } q = 3 \\ 0 & \text{otherwise} \end{cases}$$

Proof. First we note that since N is a smooth hypersurface in a weighted projective space, its cohomology is torsionfree [Dim92], and by the Lefschetz hyperplane theorem it has the cohomology of projective space for $q \neq 3$. For $q = 3$ we can compute the Alexander polynomial of the Brieskorn 7-manifold M over N by the Milnor-Orlik method described in Section 9.3 of [BG08]. This gives $H^3(M, \mathbb{Q}) = \mathbb{Q}^{42}$. But then an easy spectral sequence or Gysin sequence argument gives $H^3(N, \mathbb{Q}) \approx H^3(M, \mathbb{Q})$ which gives the result since the cohomology of N is torsionfree. \square

Now continuing Example 3.3 we have $d_0 = 0$, and $\dim_{\mathbb{C}} N = 3$. We shall assume that $d_\infty > 0$, since the no blowdown case gives rise to a join which has been studied elsewhere [BTF16, BTF22c]. Moreover, if $d_\infty > 1$ we can apply Corollary 4.3 which implies that $H^*(M, \mathbb{Q}) \approx H^*(S^{2d_\infty+3} \times N, \mathbb{Q})$. If we can take $l = 1$ in the proof of Proposition 4.1, then M will be an $S^{2d_\infty+3}$ bundle over N . So in this case when $d_\infty > 1$ the integral cohomology spectral sequence of the bundle collapses giving the cohomology groups of the product $S^{2d_\infty+3} \times N$.

As briefly indicated at the end of Section 3.1, there is a constraint on the index of N . It follows from Theorem 3.1 in [ACGTF08b] that, in order for N^{ad} to be Fano, we need the index \mathcal{J}_N of N to be at least $d_0 + 2$, so we can choose the Kähler form ω_1 on N in a way that its scalar curvature satisfies $s_1 > d_0 + 1$. Thus, we take N to be any

of the families given by 1.11, 1.12, 1.13, 1.16 in the big table of [ACC⁺21]. These are smooth hypersurfaces in weighted projective spaces. In all these cases the cohomology of N is torsionfree, and can differ from the cohomology of $\mathbb{C}\mathbb{P}^3$ only in the middle dimension. The 3rd Betti number can be computed using the Alexander polynomial as mentioned above. One can easily check in [ACC⁺21] that these projective varieties N are all polystable, and thus admit a KE metric. We indicate this in the following table of smooth Fano 3-folds.

d	\mathbf{a}	\mathbf{w}	J_N	$H^3(N, \mathbb{Z})$	KE
2	(2, 2, 2, 2, 2)	(1, 1, 1, 1, 1)	3	0	yes
3	(3, 3, 3, 3, 3)	(1, 1, 1, 1, 1)	2	\mathbb{Z}^{10}	yes
4	(4, 4, 4, 4, 2)	(1, 1, 1, 1, 2)	2	\mathbb{Z}^{20}	yes
6	(6, 6, 6, 3, 2)	(1, 1, 1, 2, 3)	2	\mathbb{Z}^{42}	yes

Remark 4.5. Applying these N to Corollary 3.5 and using Proposition 4.1 proves the existence of SE metrics on certain lens space bundles over these smooth Fano 3-folds.

APPENDIX A. CONSTANT WEIGHTED SCALAR CURVATURE ON ADMISSIBLE KÄHLER MANIFOLDS

In this appendix we will show how the technique we used in the proof of Theorem 3.1 can be adapted to an existence result for Kähler metrics of constant weighted scalar curvature. The notion of *constant weighted scalar curvature* was introduced by A. Lahdili in [Lah19].

Suppose that Ω is any admissible Kähler class on an admissible manifold N^{ad} which fibers over a local product of nonnegative CSMK metrics. We will show that when the value of the weight is sufficiently large, there exists an admissible Kähler metric in Ω with constant weighted scalar curvature. Specifically we are proving the following theorem:

Theorem A.1. *Suppose Ω is any admissible Kähler class on the admissible manifold $N^{ad} = \mathbb{P}(E_0 \oplus E_\infty) \rightarrow N$, where N is a compact Kähler manifold which is a local product of nonnegative CSMK metrics. If $p > \max\{m + 1, 2d_0 + 2, 2d_\infty + 2\}$, then there always exist some $c \in (-1, 1)$ such that the corresponding admissible $(c\mathfrak{z} + 1, p)$ -extremal metric in Ω has constant $(c\mathfrak{z} + 1, p)$ -scalar curvature.*

Note that Lemmas A.2 and A.3 below supply additional technical details for the proof of Theorem 3.1. In Section A.1 we will touch on the limitations of Theorem A.1, by exploring an example that falls outside of the realm of Theorem A.1.

Proof. Assume that we have an admissible manifold N^{ad} as above with the property that N is a local product of *non-negative* CSMK metrics and let Ω be an admissible Kähler class on N^{ad} . As we discussed in the beginning of Section 3, we know that Ω admits an admissible $(c\mathfrak{z} + 1, p)$ -extremal metric for any choice of $c \in (-1, 1)$ and $p \in \mathbb{R}$. Recall that the admissible $(c\mathfrak{z} + 1, p)$ -extremal metric has $(c\mathfrak{z} + 1, p)$ -scalar curvature $Scal_{c\mathfrak{z}+1,p} = A_1\mathfrak{z} + A_2$, where, A_1 and A_2 are given (7) and (8). Thus admissible $(c\mathfrak{z} + 1, p)$ -extremal metric has, g has constant weighted scalar curvature, $Scal_{c\mathfrak{z}+1,p}$, if and only if $A_1 = 0$. In turn the equation $A_1 = 0$ may be re-written as

$$(17) \quad \alpha_{1, -(1+p)}\beta_{0, (1-p)} - \alpha_{0, -(1+p)}\beta_{1, (1-p)} = 0.$$

As already mentioned in the proof of Theorem 3.1 we know that

- $c\alpha_{r+1,k} + \alpha_{r,k} = \alpha_{r,k+1}$
- $c\beta_{r+1,k} + \beta_{r,k} = \beta_{r,k+1}$.

Further, it is easy to check the general formulas below:

- $\frac{d}{dc} [\alpha_{r,k}] = k\alpha_{r+1,k-1}$
- $\frac{d}{dc} [\beta_{r,k}] = k\beta_{r+1,k-1}$,

These observations are used in the calculations below.

We now turn our attention to the *weighted Einstein-Hilbert functional* as defined by Futaki and Ono in [FO20]: In the present case, (the appropriate restriction of) this weighted Einstein-Hilbert functional is given by

$$WH_K(c) := \frac{S_c}{V_c^{\frac{p-2}{p}}},$$

where $V_c = \int_{N^{ad}} (c\mathfrak{z} + 1)^{-p} d\mu_g$ and $S_c = \int_{N^{ad}} \text{Scal}_{c\mathfrak{z}+1,p}(g)(c\mathfrak{z} + 1)^{-p} d\mu_g$. Using the formulas from [AMTF], it is not hard to check that, up to an overall positive rescale, $WH_K(c) = \frac{\beta_{0,2-p}}{(\alpha_{0,-p})^{\frac{p-2}{p}}}$. To simplify things a bit, we will actually work with $H_K(c) := [WH_K(c)]^p$, that is,

$$H_K(c) = \frac{(\beta_{0,2-p})^p}{(\alpha_{0,-p})^{p-2}}.$$

Note that while $H_S(c)$ from the proof of Theorem 3.1 is analogous to $H_K(c)$, it is NOT a special case of $H_K(c)$ and for $p = m + 2$, the two functions are not equal. We easily verify that

$$\begin{aligned} H'_K(c) &= \frac{p(2-p)(\alpha_{0,-p})^{p-2}(\beta_{0,2-p})^{(p-1)}\beta_{1,1-p}(-p-2)(-p)(\alpha_{0,-p})^{(p-3)}\alpha_{1,-(1+p)}(\beta_{0,2-p})^p}{(\alpha_{0,-p})^{2(p-2)}} \\ &= \frac{p(p-2)(\beta_{0,2-p})^{p-1}(\alpha_{0,-p})^{p-3}}{(\alpha_{0,-p})^{2(p-2)}} \left[\alpha_{1,-(1+p)}\beta_{0,2-p} - \alpha_{0,-p}\beta_{1,1-p} \right] \\ &= \frac{p(p-2)(\beta_{0,2-p})^{p-1}(\alpha_{0,-p})^{p-3}}{(\alpha_{0,-p})^{2(p-2)}} \left[\alpha_{1,-(1+p)}\beta_{0,(1-p)} - \alpha_{0,-(1+p)}\beta_{1,(1-p)} \right]. \end{aligned}$$

Now $\alpha_{0,-p} > 0$ and $\beta_{0,2-p} > 0$ (the latter is due to the fact that we are assuming N is a local product of *nonnegative* CSCK metrics) and therefore critical points of $H_K(c)$ correspond exactly to solutions of Equation (17). Showing that for $p > \max\{m+1, 2d_0+2, 2d_\infty+2\}$, $H_K(c)$ must have a least one critical point $c \in (-1, 1)$, would finish the proof.

To that end, first notice that $H_K(c)$ is a smooth positive function for $c \in (-1, 1)$. To show that $H(c)$ has a critical point inside $(-1, 1)$, we will show that $\lim_{c \rightarrow \pm 1^\mp} H_K(c) = +\infty$.

We need two lemmas. Let $o(x)$ denote any function which is smooth in a neighborhood of $x = 0$ and satisfies that $o(0) = 0$. We shall recycle this notation for several functions of this nature.

Lemma A.2. *For any $k \geq m + 1$,*

$$\alpha_{0,-k} = \frac{\delta_0(c) + o(1-c)}{(1-c)^{k-1-d_0}} = \frac{\delta_\infty(c) + o(1+c)}{(1+c)^{k-1-d_\infty}},$$

where δ_0 and δ_∞ are smooth functions defined in open neighborhoods of $c = 1$ and $c = -1$ respectively, such that $\delta_0(1) > 0$ and $\delta_\infty(-1) > 0$.

Proof. By repeated integration by parts, we can see that $\alpha_{0,-k} = \int_{-1}^1 (ct+1)^{-k} \overbrace{p_c(t)}^{\text{degree } m-1} dt$ is a rational function of c which is positive and has no vertical asymptotes within the interval $(-1, 1)$.

First notice that

$$\alpha_{0,-k} = \int_{-1}^1 (ct+1)^{-k} (1+t)^{d_0} p_0(t) dt = \int_{-1}^1 (ct+1)^{-k} (1-t)^{d_\infty} p_\infty(t) dt,$$

where $p_0(t) = p_c(t)/(1+t)^{d_0}$ and $p_\infty(t) = p_c(t)/(1-t)^{d_\infty}$ are polynomials of degree $m-1-d_0$ and $m-1-d_\infty$, respectively. Keep in mind that the complex dimension $m = \sum_{a \in \hat{A}} d_a + 1$ (with $d_a > 0$ for $a \in \mathcal{A}$ and $d_a \geq 0$ for $a \in \hat{\mathcal{A}}$) and note that $p_0(-1) > 0$ and $p_\infty(1) > 0$. From the definition of $p_c(t)$ we also know that $p_0(t)$ and $p_\infty(t)$ are positive for $-1 < t < 1$. To understand the behavior of $\alpha_{0,-k}$ near $c = \pm 1$, we consider repeated integration by parts and observe that

$$\alpha_{0,-k} = \frac{\left(\frac{(k-2-d_0)! d_0! p_0(-1)}{(k-1)! c^{d_0+1}} + o(1-c) \right)}{(1-c)^{k-1-d_0}} = \frac{\left(\frac{(-1)^{d_\infty+1} (k-2-d_\infty)! d_\infty! p_\infty(1)}{(k-1)! c^{d_\infty+1}} + o(1+c) \right)}{(1+c)^{k-1-d_\infty}}$$

This proves the lemma. \square

Lemma A.3. For any $l \geq m$,

$$\beta_{0,-l} = \frac{\gamma_0(c) + o(1-c)}{(1-c)^{l-d_0}} = \frac{\gamma_\infty(c) + o(1+c)}{(1+c)^{l-d_\infty}},$$

where γ_0 and γ_∞ are smooth functions defined in open neighborhoods of $c = 1$ and $c = -1$ respectively, such that $\gamma_0(1) > 0$ and $\gamma_\infty(-1) > 0$.

Proof. By repeated integration by parts, we can see that

$$\begin{aligned} \beta_{0,-l} &= \int_{-1}^1 \overbrace{\left(\sum_{a \in \hat{A}} \frac{x_a d_a s_a}{1+x_a t} \right)}^{\text{degree } m-2} p_c(t) (ct+1)^{-l} dt \\ &\quad + \left((1-c)^{-l} p_c(-1) + (1+c)^{-l} p_c(1) \right) \end{aligned}$$

is a rational function of c which is non-negative and has no vertical asymptotes within the interval $(-1, 1)$. Moreover, it is easy to see that if $d_0 = 0$,

$$\beta_{0,-l} = \frac{(p_c(-1) + o(1-c))}{(1-c)^l},$$

with $p_c(-1) > 0$ in this case, and if $d_\infty = 0$,

$$\beta_{0,-l} = \frac{(p_c(1) + o(1+c))}{(1-c)^l}$$

with $p_c(1) > 0$ in this case.

Now assume that $d_0 \geq 1$ and define

$$q_0(t) := \sum_{a \in \hat{A} \setminus \{0\}} \frac{x_a d_a s_a}{1 + x_a t} \frac{p_c(t)}{(1+t)^{d_0-1}} + x_0 d_0 s_0 \frac{p_c(t)}{(1+t)^{d_0}}.$$

The degree of the polynomial $q_0(t)$ is $m - d_0 - 1$ and

$$q_0(-1) = x_0 d_0 s_0 \prod_{a \in \hat{A} \setminus \{0\}} (1 - x_a)^{d_a} = d_0(d_0 + 1) \prod_{a \in \hat{A} \setminus \{0\}} (1 - x_a)^{d_a} > 0.$$

By repeated integration by parts we then have

$$\begin{aligned} \beta_{0,-l} &= \int_{-1}^1 (ct + 1)^{-l} (1+t)^{d_0-1} q_0(t) dt + (1+c)^{-l} p_c(1) \\ &= \frac{\left(\frac{(l-2-(d_0-1))! (d_0-1)! q_0(-1)}{(l-1)! c^{d_0}} + o(1-c) \right)}{(1-c)^{l-d_0}}. \end{aligned}$$

Similarly if $d_\infty \geq 1$ we define

$$q_\infty(t) := \sum_{a \in \hat{A} \setminus \{\infty\}} \frac{x_a d_a s_a}{1 + x_a t} \frac{p_c(t)}{(1-t)^{d_\infty-1}} + x_\infty d_\infty s_\infty \frac{p_c(t)}{(1-t)^{d_\infty}}.$$

The degree of the polynomial $q_\infty(t)$ is $m - d_\infty - 1$ and

$$q_\infty(1) = x_\infty d_\infty s_\infty \prod_{a \in \hat{A} \setminus \{\infty\}} (1 + x_a)^{d_a} = d_\infty(d_\infty + 1) \prod_{a \in \hat{A} \setminus \{\infty\}} (1 + x_a)^{d_a} > 0.$$

By repeated integration by parts we now have

$$\begin{aligned} \beta_{0,-l} &= \int_{-1}^1 (ct + 1)^{-l} (1-t)^{d_\infty-1} q_\infty(t) dt + (1-c)^{-l} p_c(-1) \\ &= \frac{\left(\frac{(-1)^{d_\infty} (l-2-(d_\infty-1))! (d_\infty-1)! q_\infty(1)}{(l-1)! c^{d_\infty}} + o(c+1) \right)}{(c+1)^{l-d_\infty}}. \end{aligned}$$

From the above observations we have now proved Lemma A.3 regardless of the values of d_0 and d_∞ . \square

From Lemmas A.2 and A.3 we now have that for $k \geq m + 1$ and $l \geq m$,

$$\frac{(\beta_{0,-l})^k}{(\alpha_{0,-k})^l} = \frac{(\gamma_0(c) + o(1-c))^k}{(\delta_0(c) + o(1-c))^l} (1-c)^{d_0(k-l)-l} = \frac{(\gamma_\infty(c) + o(1+c))^k}{(\delta_\infty(c) + o(1+c))^l} (1+c)^{d_\infty(k-l)-l}.$$

In particular, for $p \geq m + 2$, we see that (letting $l = p - 2$ and $k = p$),

$$H_K(c) = \frac{(\beta_{0,2-p})^p}{(\alpha_{0,-p})^{p-2}} = \frac{(\gamma_0(c) + o(1-c))^p}{(\delta_0(c) + o(1-c))^{p-2}} (1-c)^{2d_0+2-p} = \frac{(\gamma_\infty(c) + o(1+c))^p}{(\delta_\infty(c) + o(1+c))^{p-2}} (1+c)^{2d_\infty+2-p}.$$

Therefore, for $p > \max\{m + 1, 2d_0 + 2, 2d_\infty + 2\}$, $\lim_{c \rightarrow \pm 1^\mp} H_K(c) = +\infty$. This, together with the fact that $H_K(c)$ is smooth and bounded from below over the interval $(-1, 1)$, allows us to conclude that $H_K(c)$ has at least one critical point $c \in (-1, 1)$. For this value of c , we have $H'_K(c) = 0$ and therefore c solves (17). \square

Recall that $m = \sum_{a \in \hat{A}} d_a + 1$. Assuming (as we should) that $\mathcal{A} \neq \emptyset$, we have that for $p = 2m$ the condition $p > \max\{m + 1, 2d_0 + 2, 2d_\infty + 2\}$ is automatic. As a special case, we thus get the following corollary to Theorem A.1, which in particular confirms Conjecture 1 in [AMTF]. Note that the corollary is also a special case of an existence theorem stated in [Gua23].

Corollary A.4. *Suppose Ω is any admissible Kähler class on the admissible manifold $N^{ad} = \mathbb{P}(E_0 \oplus E_\infty) \rightarrow N$, where N is a compact Kähler manifold which is a local product of nonnegative CSCK metrics. Then there always exist some $c \in (-1, 1)$ such that the corresponding admissible $(c\mathfrak{z} + 1, 2m)$ -extremal metric g in Ω has the property that $h = (c\mathfrak{z} + 1)^{-2}g$ has constant scalar curvature and therefore is conformally Kähler Einstein-Marwell.*

A.1. Limitation of Theorem A.1. On a more pessimistic note, if we instead work with $p = m + 2$, we cannot assume that $p > \max\{m + 1, 2d_0 + 2, 2d_\infty + 2\}$ is true and thus we cannot appeal to Theorem A.1. The following example illustrates that, in this case, existence is more unpredictable - even if the underlying admissible manifold is fixed.

Example A.1. Let $\mathcal{A} = \{1\}$, $N = N_1 = \mathbb{C}\mathbb{P}^1$, and $s_1 = 2$ (so g_1 is the Fubini-Study metric on $\mathbb{C}\mathbb{P}^1$). If we then let $d_\infty = 1$ and $d_0 = 3$, we have $m = 6$ and $p = 8$. Clearly $p = 2d_0 + 2$ and hence $p > \max\{m + 1, 2d_0 + 2, 2d_\infty + 2\}$ is false. Setting $x_1 = x$ we have that each value $0 < x < 1$ determines a certain admissible Kähler class. We will see how the existence of a solution $c \in (-1, 1)$ to (17) depends on the value of x .

We calculate that

$$\begin{aligned} f_x(c) &:= \alpha_{1,-9}\beta_{0,-7} - \alpha_{0,-9}\beta_{1,-7} \\ &= -12x + (24 + x + 13x^2)c + (-17 - 52x + x^2)c^2 + (-1 + 15x + 28x^2)c^3. \end{aligned}$$

Thus solutions $c \in (-1, 1)$ to (17) correspond to roots in $(-1, 1)$ of the cubic $f_x(c)$. This cubic (with real coefficients) has discriminant D_x given by

$$D_x = -5(1+x)^3(-44352 + 155904x - 159125x^2 + 115249x^3 - 90591x^4 + 49179x^5).$$

We can check numerically that there exists a specific value $\tilde{x} \in (0, 1)$ such that $D_{\tilde{x}} = 0$ and for $0 < x < \tilde{x}$, $D_x > 0$, while for $\tilde{x} < x < 1$, $D_x < 0$. Note that $\tilde{x} \approx 0.429$.

For $\tilde{x} < x < 1$, we know that the cubic $f_x(c)$ has exactly one real root \hat{c} . One can check that for this range of x values, $f_x(1) < 1$ and $\lim_{c \rightarrow +\infty} f_x(c) = +\infty$, so it is clear that $\hat{c} > 1$ and thus for $\tilde{x} < x < 1$, $f_x(c)$ has no roots in $(-1, 1)$. Therefore, the admissible Kähler classes corresponding to values $\tilde{x} < x < 1$ admit no admissible Kähler metrics with constant $(c\mathfrak{z} + 1, 8)$ -scalar curvature. The question remains whether there could be any Kähler metric in those Kähler classes with constant $(f, 8)$ -scalar curvature, where f is some positive killing potential.

For $0 < x < \tilde{x}$, we know that the cubic $f_x(c)$ has three distinct real roots. We will see that at least one of them will be inside $(-1, 1)$.

- In the case where $0 < x < 1/7$, this is completely straightforward since $f_x(-1) = -40(1+x)^2 < 0$ and $f_x(1) = 6(1-x)(1-7x) > 0$.

- In the case where $x = 1/7$, we have $f_{1/7}(c) = \frac{4}{49}(c-1)(21c^2 - 278c + 21)$ and it is easy to check that $21c^2 - 278c + 21$ has a root in $(-1, 1)$.
- In the case where $1/7 < x < \tilde{x}$, we have $f_x(-1) < 0$ and $f_x(1) < 0$, but we also have that $f'_x(-1) = 5(1+x)(11+19x) > 0$, $f'_x(1) = -13 - 58x + 99x^2 < 0$, and $\lim_{c \rightarrow +\infty} f_x(c) = +\infty$. Considering the options for the cubic with three distinct real roots, it is clear that two of the roots must be inside $(-1, 1)$ and that the third root is in the interval $(1, +\infty)$.

Finally, for $x = \tilde{x}$, we know that the cubic $f_x(c)$ has a multiple root (and all the roots are real). In this case we still have that $f_{\tilde{x}}(-1) < 0$, $f'_{\tilde{x}}(-1) > 0$, $f_{\tilde{x}}(1) < 0$, $f'_{\tilde{x}}(1) < 0$, and $\lim_{c \rightarrow +\infty} f_x(c) = +\infty$. It is therefore clear that in this case $f_{\tilde{x}}(c)$ has a double real root in the interval $(-1, 1)$ and a single real root in the interval $(1, +\infty)$. In conclusion, for each admissible Kähler class, Ω_x , corresponding to a value $0 < x \leq \tilde{x}$, there exists at least one value $c \in (-1, 1)$ such that Ω_x admits an admissible Kähler metric with constant $(c\mathfrak{J} + 1, 8)$ -scalar curvature.

REFERENCES

- [AC21] Vestislav Apostolov and David M. J. Calderbank, *The CR geometry of weighted extremal Kähler and Sasaki metrics*, Math. Ann. **379** (2021), no. 3-4, 1047–1088. MR 4238260
- [ACC⁺21] Carolina Araujo, Ana-Maria Castravet, Ivan Cheltsov, Kento Fujita, Anne-Sophie Kaloghiros, Jesus Martinez-Garcia, Constantin Shramov, Hendrik Süß, and Nivedita Viswanathan, *The Calabi problem for Fano threefolds*, workingpaper 31, 2021.
- [ACGTF08a] Vestislav Apostolov, David M. J. Calderbank, Paul Gauduchon, and Christina W. Tønnesen-Friedman, *Hamiltonian 2-forms in Kähler geometry. III. Extremal metrics and stability*, Invent. Math. **173** (2008), no. 3, 547–601. MR MR2425136 (2009m:32043)
- [ACGTF08b] ———, *Hamiltonian 2-forms in Kähler geometry. IV. Weakly Bochner-flat Kähler manifolds*, Comm. Anal. Geom. **16** (2008), no. 1, 91–126. MR 2411469 (2010c:32043)
- [ACL21] Vestislav Apostolov, David M. J. Calderbank, and Eveline Legendre, *Weighted K-stability of polarized varieties and extremality of Sasaki manifolds*, Adv. Math. **391** (2021), Paper No. 107969, 63. MR 4300921
- [AJL] Vestislav Apostolov, Simon Jubert, and Abdellah Lohdili, *Weighted K-Stability and coercivity with applications to Extremal Kähler and Sasaki Metrics*, arXiv: 2104.09709v2, to appear in Geometry & Topology.
- [AMTF] Vestislav Apostolov, Gideon Maschler, and Christina W. Tønnesen-Friedman, *Weighted extremal Kähler metrics and the Einstein-Maxwell geometry of projective bundles*, arXiv: 1808.02813 (to appear in Comm. Anal. Geom.).
- [AZ21] Hamid Ahmadinezhad and Ziquan Zhuang, *Seshadri constants and K-stability of Fano manifolds*, arXiv:2101.09246 (2021).
- [BCTF19] Charles P. Boyer, David M. J. Calderbank, and Christina W. Tønnesen-Friedman, *The Kähler geometry of Bott manifolds*, Adv. Math. **350** (2019), 1–62. MR 3945589
- [BG08] Charles P. Boyer and Krzysztof Galicki, *Sasakian geometry*, Oxford Mathematical Monographs, Oxford University Press, Oxford, 2008. MR MR2382957 (2009c:53058)
- [BGK05] C. P. Boyer, K. Galicki, and J. Kollár, *Einstein metrics on spheres*, Ann. of Math. (2) **162** (2005), no. 1, 557–580. MR 2178969 (2006j:53058)
- [BGO07] Charles P. Boyer, Krzysztof Galicki, and Liviu Ornea, *Constructions in Sasakian geometry*, Math. Z. **257** (2007), no. 4, 907–924. MR MR2342558 (2008m:53103)
- [BGS08] Charles P. Boyer, Krzysztof Galicki, and Santiago R. Simanca, *Canonical Sasakian metrics*, Commun. Math. Phys. **279** (2008), no. 3, 705–733. MR MR2386725

- [BHL18] Charles Boyer, Hongnian Huang, and Eveline Legendre, *An application of the Duistermaat–Heckman theorem and its extensions in Sasaki geometry*, *Geom. Topol.* **22** (2018), no. 7, 4205–4234. MR 3890775
- [BHLTF17] Charles P. Boyer, Hongnian Huang, Eveline Legendre, and Christina W. Tønnesen-Friedman, *The Einstein–Hilbert functional and the Sasaki–Futaki invariant*, *Int. Math. Res. Not. IMRN* (2017), no. 7, 1942–1974. MR 3658189
- [BHLTF18] ———, *Reducibility in Sasakian geometry*, *Trans. Amer. Math. Soc.* **370** (2018), no. 10, 6825–6869. MR 3841834
- [BHLTF21] ———, *Some open problems in Sasaki geometry*, *Differential geometry in the large*, *London Math. Soc. Lecture Note Ser.*, vol. 463, Cambridge Univ. Press, Cambridge, 2021, pp. 143–168. MR 4420789
- [Boy13] Charles P. Boyer, *Maximal tori in contactomorphism groups*, *Differential Geom. Appl.* **31** (2013), no. 2, 190–216. MR 3032643
- [BT82] R. Bott and L. W. Tu, *Differential forms in algebraic topology*, *Graduate Texts in Mathematics*, vol. 82, Springer-Verlag, New York, 1982. MR 658304 (83i:57016)
- [BTF13] Charles P. Boyer and Christina W. Tønnesen-Friedman, *Extremal Sasakian geometry on $T^2 \times S^3$ and related manifolds*, *Compos. Math.* **149** (2013), no. 8, 1431–1456. MR 3103072
- [BTF14] ———, *Extremal Sasakian geometry on S^3 -bundles over Riemann surfaces*, *Int. Math. Res. Not. IMRN* (2014), no. 20, 5510–5562. MR 3271180
- [BTF16] ———, *The Sasaki join, Hamiltonian 2-forms, and constant scalar curvature*, *J. Geom. Anal.* **26** (2016), no. 2, 1023–1060. MR 3472828
- [BTF21] ———, *Sasakian geometry on sphere bundles*, *Differential Geom. Appl.* **77** (2021), Paper No. 101765, 37. MR 4253896
- [BTF22a] ———, *Constant scalar curvature Sasaki metrics and projective bundles*, arXiv:2112.01567, to appear in *Birational Geometry, Kähler–Einstein Metrics, and Degeneration in the Springer Proceedings in Mathematics & Statistics*. (2022).
- [BTF22b] ———, *Iterated S^3 Sasaki joins and Bott orbifolds*, *Ann. Fac. Sci. Toulouse Math.* (6) **31** (2022), no. 3, 837–860. MR 4452254
- [BTF22c] ———, *The S^3 Sasaki join construction*, arXiv:1911.11031, to appear in the *J. Math. Soc. Japan* (2022).
- [BvC18] Charles P. Boyer and Craig van Coevering, *Relative K -stability and extremal Sasaki metrics*, *Math. Res. Lett.* **25** (2018), no. 1, 1–19. MR 3818612
- [Cal82] E. Calabi, *Extremal Kähler metrics*, *Seminar on Differential Geometry*, *Ann. of Math. Stud.*, vol. 102, Princeton Univ. Press, Princeton, N.J., 1982, pp. 259–290. MR 83i:53088
- [CS09] I. A. Cheltsov and K. A. Shramov, *Extremal metrics on del Pezzo threefolds*, *Tr. Mat. Inst. Steklova* **264** (2009), no. Mnogomernaya Algebraicheskaya Geometriya, 37–51. MR 2590832
- [CS18] Tristan C. Collins and Gábor Székelyhidi, *K -semistability for irregular Sasakian manifolds*, *J. Differential Geom.* **109** (2018), no. 1, 81–109. MR 3798716
- [Dim92] A. Dimca, *Singularities and topology of hypersurfaces*, *Universitext*, Springer-Verlag, New York, 1992. MR 94b:32058
- [FO20] Akito Futaki and Hajime Ono, *On the existence problem of Einstein–Maxwell Kähler metrics*, *Geometric analysis—in honor of Gang Tian’s 60th birthday*, *Progr. Math.*, vol. 333, Birkhäuser/Springer, Cham, [2020] ©2020, pp. 93–111. MR 4180998
- [FOW09] Akito Futaki, Hajime Ono, and Guofang Wang, *Transverse Kähler geometry of Sasaki manifolds and toric Sasaki–Einstein manifolds*, *J. Differential Geom.* **83** (2009), no. 3, 585–635. MR MR2581358
- [Gua95] Daniel Guan, *Existence of extremal metrics on compact almost homogeneous Kähler manifolds with two ends*, *Trans. Amer. Math. Soc.* **347** (1995), no. 6, 2255–2262. MR 1285992 (96a:58059)
- [Gua23] Zhuangdan Guan, *Maxwell–Einstein metrics on completions of certain C^* bundles*, *Acta Math. Sci.* **43** (2023), 363–372.

- [HS02] Andrew D. Hwang and Michael A. Singer, *A momentum construction for circle-invariant Kähler metrics*, Trans. Amer. Math. Soc. **354** (2002), no. 6, 2285–2325 (electronic). MR 1885653 (2002m:53057)
- [HS16] Weiyong He and Song Sun, *Frankel conjecture and Sasaki geometry*, Adv. Math. **291** (2016), 912–960. MR 3459033
- [Hwa94] Andrew D. Hwang, *On existence of Kähler metrics with constant scalar curvature*, Osaka J. Math. **31** (1994), no. 3, 561–595. MR 1309403 (96a:53061)
- [Koi90] Norihito Koiso, *On rotationally symmetric Hamilton's equation for Kähler-Einstein metrics*, Recent topics in differential and analytic geometry, Adv. Stud. Pure Math., vol. 18, Academic Press, Boston, MA, 1990, pp. 327–337. MR 1145263 (93d:53057)
- [KS86] Norihito Koiso and Yusuke Sakane, *Nonhomogeneous Kähler-Einstein metrics on compact complex manifolds*, Curvature and topology of Riemannian manifolds (Katata, 1985), Lecture Notes in Math., vol. 1201, Springer, Berlin, 1986, pp. 165–179. MR 859583 (88c:53047)
- [KT91] Matthias Kreck and Georgia Triantafyllou, *On the classification of manifolds up to finite ambiguity*, Canad. J. Math. **43** (1991), no. 2, 356–370. MR 1113760 (92h:57041)
- [Lah19] Abdellah Lahdili, *Kähler metrics with constant weighted scalar curvature and weighted K -stability*, Proc. Lond. Math. Soc. (3) **119** (2019), no. 4, 1065–1114. MR 3964827
- [LeB91] C. LeBrun, *Scalar-flat Kähler metrics on blown-up ruled surfaces*, J. Reine Angew. Math. **420** (1991), 161–177. MR 1124569 (92i:53066)
- [Ler02] E. Lerman, *A convexity theorem for torus actions on contact manifolds*, Illinois J. Math. **46** (2002), no. 1, 171–184. MR 2003g:53146
- [MN13] Toshiki Mabuchi and Yasuhiro Nakagawa, *New examples of Sasaki-Einstein manifolds*, Tohoku Math. J. (2) **65** (2013), no. 2, 243–252. MR 3079287
- [MTF11] Gideon Maschler and Christina W. Tønnesen-Friedman, *Generalizations of Kähler-Ricci solitons on projective bundles*, Math. Scand. **108** (2011), no. 2, 161–176. MR 2805600
- [PP98] H. Pedersen and Y.-S. Poon, *Deformations of hypercomplex structures*, J. Reine Angew. Math. **499** (1998), 81–99. MR 99h:32009
- [Sim91] S. R. Simanca, *Kähler metrics of constant scalar curvature on bundles over $\mathbb{C}P_{n-1}$* , Math. Ann. **291** (1991), no. 2, 239–246. MR 1129363 (92m:53115)
- [TF98] Christina Wiis Tønnesen-Friedman, *Extremal Kähler metrics on minimal ruled surfaces*, J. Reine Angew. Math. **502** (1998), 175–197. MR MR1647571 (99g:58026)
- [Yam99] T. Yamazaki, *A construction of K -contact manifolds by a fiber join*, Tohoku Math. J. (2) **51** (1999), no. 4, 433–446. MR 2001e:53094

CHARLES P. BOYER, DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF NEW MEXICO, ALBUQUERQUE, NM 87131.

Email address: cboyer@unm.edu

HONGNIAN HUANG, DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF NEW MEXICO, ALBUQUERQUE, NM 87131.

Email address: hnhuang@gmail.com

EVELINE LEGENDRE, UNIVERSITÉ CLAUDE BERNARD (LYON 1), INSTITUT CAMILLE JORDAN, 43 BOULEVARD DU 11 NOVEMBRE 1918 F-69622 VILLEURBANNE CEDEX, FRANCE

Email address: legendre@math.univ-lyon1.fr

CHRISTINA W. TØNNESEN-FRIEDMAN, DEPARTMENT OF MATHEMATICS, UNION COLLEGE, SCHENECTADY, NEW YORK 12308, USA

Email address: tonnesec@union.edu